arXiv:1101.4971v2 [math.GT] 22 Mar 2011

THE GEOMETRY OF CYCLIC HYPERBOLIC POLYGONS

JASON DEBLOIS

Abstract. A polygon in the hyperbolic plane is *cyclic* if a single circle contains all of its vertices; we will say it is *centered* if in addition its interior contains the center of this circle. We give necessary and sufficient conditions for a set of real numbers to be the side length collection of a cyclic or centered polygon. A cyclic polygon is uniquely determined by its collection of side lengths; its vertex angles vary as C^1 functions of side lengths; and so does the radius of the circle containing its vertices. We describe the derivatives of these quantities and show in consequence that the area of centered polygons is "monotonic" in side lengths. This does not hold for non-centered cyclic polygons, but for these the *horocyclic* polygons, those with all vertices on a horocycle, can be used to bound area below.

J.M. Schlenker showed that an n-tuple $(d_0, \ldots, d_{n-1})$ of positive real numbers is the side length collection of an n-gon P in $\mathbb{H}^2$ that is cyclic or horocyclic, or has all vertices equidistant from a fixed geodesic, as long as $d_i \leq \sum_{j\neq i} d_j$ for each i [1]; and furthermore, that P is uniquely prescribed by (the equivalence class, under cyclic permutation of entries, of) this n-tuple, up to orientation-preserving isometry of $\mathbb{H}^2$. We will show that P is cyclic if and only if for each i,

$$\sinh(d_i/2) < \sum_{j\neq i} \sinh(d_j/2);$$

and P is horocyclic if and only if there is some i such that equality holds above. This fact may also be implicit in a recent preprint of R. Walter [2] that re-proves Schlenker's result. It serves as the starting point for our main aim here: to describe how certain geometric quantities vary over the entire set of cyclic n-gons.

Given the fact above it is natural to parametrize the collection of (marked) cyclic n-gons in $\mathbb{H}^2$ by a subset $\widetilde{\mathcal{AC}}_n$ of $\mathbb{R}^n$ whose points correspond to side length collections. With this perspective we will show that the area of a cyclic polygon, and the radius of the circle containing its vertices, determine continuously differentiable functions on $\widetilde{\mathcal{AC}}_n$; we will give formulas and bounds for their derivatives; and we will bound the area of a cyclic polygon below given bounds on its side lengths.

A key theme is that area displays qualitatively different behaviors on $\widetilde{\mathcal{C}}_n$, the set of centered n-gons, versus $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$: increasing any side length of a centered n-gon increases its area, and this also holds for all sides of a non-centered n-gon but the longest; but increasing the largest side length of a non-centered cyclic n-gon decreases its area. The set $\widetilde{\mathcal{HC}}_n$ of horocyclic n-gons, parametrized by side length, is the frontier of $\widetilde{\mathcal{AC}}_n$ in $\mathbb{R}^n$. We will show that area extends continuously to $\widetilde{\mathcal{HC}}_n$, and use its values here to produce lower bounds.

Partially supported by NSF grant DMS-1007175.

The paper is structured as follows. Section 1 introduces cyclic and centered polygons and proves some basic results about their structure. Most importantly, Lemma 1.6 describes an "isosceles decomposition," and Lemma 1.10 uses this to produce two equations: one relates the side lengths of a centered polygon P to the radius of the circle containing its vertices, and the other applies if P is cyclic but not centered.

The main result of Section 2, Proposition 2.7 gives necessary and sufficient conditions for an n-tuple of positive real numbers to be the side length collection of a cyclic or centered n-gon in $\mathbb{H}^2$, and also asserts uniqueness, up to isometry of n-gons and cyclic permutation of entries. Proposition 2.9 is the analogous result for horocyclic n-gons.

Definition 3.1 parametrizes the sets $\widetilde{\mathcal{AC}}_n \subset \mathbb{R}^n$, of marked cyclic n-gons, and $\widetilde{\mathcal{C}}_n \subset \widetilde{\mathcal{AC}}_n$, of centered n-gons, by their side length collections. We then re-interpret the radius of the circle containing the endpoints of a cyclic n-gon and certain relevant angle measures as values of functions on $\widetilde{\mathcal{AC}}_n$, and we describe equivariance properties under the action of cyclic permutation of entries. In Section 3 we also describe the frontier of $\widetilde{\mathcal{C}}_n$ in $\widetilde{\mathcal{AC}}_n$ (Lemma 3.3) and show that the set $\widetilde{\mathcal{HC}}_n$ of horocyclic n-gons is the frontier of $\widetilde{\mathcal{AC}}_n$ in $\mathbb{R}^n$ (Lemma 3.10).

Proposition 4.1 asserts in particular that the radius of cyclic n-gons depends in continuously differentiable fashion on their side lengths, and Proposition 4.4 confirms the same for the angle at a vertex. We also describe the limiting behavior of radius and vertex angle as polygons approach horocyclic, in Lemma 4.7 and Corollary 4.9, respectively.

Section 5 considers the area of a cyclic polygon P, or more generally, the *radius-R defect* of P. For certain $R \geq 0$ and $P \in \widetilde{\mathcal{C}}_n$, this is the area of the region in P complementary to a collection of disjoint disks with radius R centered at its vertices. The derivative of this quantity is described in Proposition 5.5, from which it follows that the defect of centered polygons is "monotonic" in the side lengths (Corollary 5.8). As mentioned above, this does not hold on $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$, but Corollary 5.11 gives bounds using the defect's extension to $\widetilde{\mathcal{HC}}_n$.

Section 6 collects some computationally useful explicit formulas. Notably, Lemma 6.1 gives one for the radius of a cyclic triangle, using hyperbolic trigonometric functions of side lengths.

We are motivated here by an application to Voronoi and Delaunay tessellations of hyperbolic surfaces, to be explored in a later paper. Our methods are elementary: we use only standard facts of hyperbolic geometry, some basic calculus and differential topology, and at one point the Jordan curve theorem. Sorry about the length; we hope it is at least a decent read.

1. Basic structure of cyclic and centered polygons

In the hyperbolic plane $\mathbb{H}^2$, a *half-plane* bounded by a geodesic γ is the closure of a component of $\mathbb{H}^2 - \gamma$. A *convex polygon* is the nonempty intersection of a collection of half-planes $\{\mathcal{H}_i\}$ such that the associated collection of frontier geodesics $\{\gamma_i\}$ has the property that $\#\{i \,|\, \gamma_i \cap B_r(p) \neq \emptyset\} < \infty$ for each $r > 0$ and $p \in \mathbb{H}^2$, where $B_r(p) = \{x \in \mathbb{H}^2 \,|\, d(x,p) < r\}$ is the open disk of radius r centered at p. We will use the following standard facts repeatedly:

Facts. Given $p \in \mathbb{H}^2$:



FIGURE 1.1. In bold: a triangle T that is centered, and one that is cyclic but not centered, each with center v and radius J.

- For any one-dimensional subspace V of $T_p\mathbb{H}^2$, there is a unique geodesic line γ in $\mathbb{H}^2$ with $p \in \gamma$ and $V = T_p\gamma$.
- For any $q \neq p$ there is a unique geodesic arc γ with endpoints p and q, and a unique geodesic line δ in $\mathbb{H}^2$ with $\gamma \subset \delta$.

These imply for example that if non-identical geodesics intersect, they do so transversely and at a single point. They also imply that half-planes are *convex*; that is, for p and q in a half-plane $\mathcal{H}$, the unique geodesic arc joining p to q is entirely contained in $\mathcal{H}$. The intersection of convex sets is clearly convex, so a "convex polygon," as we defined it, is convex. (There is a weaker definition of "polygon" than ours, that does not imply convexity.)

An *edge* (or *side*) of $P = \bigcap_i \mathcal{H}_i$ is $\gamma_i \cap P$ for some i such that this intersection is not empty or a singleton. A *vertex* of P is the nonempty intersection of two edges. The *boundary* ∂P of P is the union of its edges. It is easy to see that ∂P is the topological frontier $P \cap \overline{\mathbb{H}^2 - P}$ of P in $\mathbb{H}^2$ and has the structure of a *Jordan curve*, the image of an embedding of S^1.

Definition 1.1. A *circle* in $\mathbb{H}^2$ is the set of points at fixed distance (its *radius*) from a fixed point (its *center*). A compact, convex polygon $P \subset \mathbb{H}^2$ is *cyclic* if there is a circle in $\mathbb{H}^2$ containing all of its vertices and *centered* if, additionally, the center of this circle is contained in the interior of P (see Figure 1.1). We refer by the *center* and *radius* of a cyclic polygon to the center and radius, respectively, of the circle containing its vertices.

A cyclic n-gon P uniquely determines its center, since this is the intersection of the perpendicular bisectors to any two edges of P, and hence also its radius.

Definition 1.2. If half-planes $\mathcal{H}$ and $\mathcal{H}'$ have frontier geodesics γ and γ', respectively, that meet at a point p, the *interior angle at* p between $\mathcal{H}$ and $\mathcal{H}'$ is the value $\theta \in [0, \pi]$ such that $\langle \eta, \eta' \rangle = \cos(\pi - \theta)$, where η and η' are outward-pointing unit normal vectors in $T_p\mathbb{H}^2$ to $\mathcal{H}$ and $\mathcal{H}'$, respectively, and $\langle , \rangle$ is the hyperbolic inner product.

If P is a convex polygon with edges e and e' intersecting at a vertex v, the *angle of* P *at* v, denoted $\angle_v P$, is the interior angle at v between $\mathcal{H}$ and $\mathcal{H}'$, where $\mathcal{H}$ (or respectively, $\mathcal{H}'$) is the half-plane containing P with frontier geodesic containing e (resp. e').

The facts previously stated imply that with $\mathcal{H}$, $\mathcal{H}'$ and p as above the interior angle at p between $\mathcal{H}$ and $\mathcal{H}'$ is equal to π if and only if $\mathcal{H} = \mathcal{H}'$ and 0 if and only if $\mathcal{H}$ and $\mathcal{H}'$ are *opposite*; that is, there is a geodesic γ such that $\mathcal{H}$ is the closure of one component of $\mathbb{H}^2 - \gamma$, and $\mathcal{H}'$ is the closure of the other.

We will use the *hyperbolic law of cosines* extensively below: Let P be a hyperbolic triangle with edges $\gamma_0, \gamma_1, \gamma_2$ of length d_0, d_1, d_2, respectively. If α_i is the angle of P at the vertex opposite γ_i ($i \in \{0,1,2\}$), then:

$$\cosh d_2 = \cosh d_0 \cosh d_1 - \sinh d_0 \sinh d_1 \cos \alpha_2, \tag{1.2.1}$$

The *hyperbolic law of sines* asserts:

$$\frac{\sin \alpha_0}{\sinh d_0} = \frac{\sin \alpha_1}{\sinh d_1} = \frac{\sin \alpha_2}{\sinh d_2} \tag{1.2.2}$$

These imply that for any $r > 0$ and $p \in \mathbb{H}^2$, the open disk $B_r(p) = \{x \in \mathbb{H}^2 \mid d(x,p) < r\}$ is convex; so is its closure $\overline{B}_r(p) = \{x \in \mathbb{H}^2 \mid d(x,p) \le r\}$; and that no three points on the circle $C = \partial B_r(p)$ lie on the same hyperbolic geodesic.

Definition 1.3. A finite collection $\{x_0, x_1, \ldots, x_{n-1}\}$ of distinct points on a circle $C \subset \mathbb{H}^2$ is *cyclically ordered* if for each i a single component of $C - \{x_i, x_{i-1}\}$ contains x_j for all $j \neq i$ or $i-1 \,(\mathrm{modulo}\, n)$.

Lemma 1.4. *Given a circle $C \subset \mathbb{H}^2$ and $\mathcal{S} = \{x_0, x_1, \ldots, x_{n-1}\} \subset C$ cyclically ordered, for each i let γ_i be the geodesic arc joining x_{i-1} to x_i, taking $i-1$ modulo n, and let δ_i be the geodesic in $\mathbb{H}^2$ containing γ_i. There is a unique cyclic n-gon $P = \bigcap_{i=0}^{n-1} \mathcal{H}_i$ with vertex set $\mathcal{S}$ and edge set $\{\gamma_i\}$, where for each i, $\mathcal{H}_i$ is a half space bounded by δ_i.*

Proof. For $j \neq i$ or $i \pm 1 \,(\mathrm{mod}\, n)$, since the endpoints of γ_j do not separate those of γ_i on C it follows that $\gamma_i \cap \gamma_j = \emptyset$. Therefore $\bigcup_{i=0}^{n-1} \gamma_i$ has the structure of a Jordan curve in $\mathbb{H}^2$, so the Jordan curve theorem implies that it separates $\mathbb{H}^2$ into two components: one with compact closure and one without. Let P^o be the component of $\mathbb{H}^2 - \bigcup_{i=0}^{n-1} \gamma_i$ with compact closure, and let P be its closure. Then $\partial P \doteq \bigcup_{i=0}^{n-1} \gamma_i$ is the topological frontier of P and P^o in $\mathbb{H}^2$. We will show that P is a convex polygon.

By definition C is the frontier of the open disk $B_J(v)$ of radius J, where v is the center and J the radius of C. Since $\overline{B}_J(v)$ is convex and contains the endpoints of γ_i for each i, it also contains γ_i, and hence $\partial P \subset \overline{B}_J(v)$. It follows that $P \subset \overline{B}_J(v)$ and $P^o \subset B_J(v)$. Since $\overline{B}_J(v)$ is convex, $\gamma_i = \delta_i \cap \overline{B_J(v)}$. Since P^o is a connected open set contained in $B_J(v) - \gamma_i$, it does not intersect δ_i and hence lies in one component of $\mathbb{H}^2 - \delta_i$. Let $\mathcal{H}_i$ be the closure of this component; then $P \subset \mathcal{H}_i$ and $\partial P \cap \partial \mathcal{H}_i = \gamma_i$.

Let δ_i^+ be the component of $\delta_i - \gamma_i$ with one endpoint at x_i, and let δ_i^- be the component with an endpoint at x_{i-1}. Since $\delta_i \cap \delta_{i+1} = \gamma_i \cap \gamma_{i+1} = \{x_i\}$, and $\gamma_i \subset P \subset \mathcal{H}_{i+1}$, the component of $\delta_i - \{x_i\}$ that does not intersect $\mathcal{H}_{i+1}$ is δ_i^+. Similarly δ_i^- does not intersect $\mathcal{H}_{i-1}$. (Above, take $i \pm 1$ modulo n when necessary.)

By construction, P is contained in $\bigcap_{i=0}^{n-1} \mathcal{H}_i$, and the frontier of $\bigcap_{i=0}^{n-1} \mathcal{H}_i$ is contained in $\bigcup_{i=0}^{n-1} \left(\delta_i - (\delta_i^- \cup \delta_i^+)\right) = \bigcup_{i=0}^{n-1} \gamma_i$. Therefore $P = \bigcap_{i=0}^{n-1} \mathcal{H}_i$ is a convex polygon with edge set $\{\gamma_i\}$ and vertex set $\mathcal{S}$. Uniqueness also follows by construction. $\square$

The next lemma records an important observation about non-centered cyclic polygons.

Lemma 1.5. *Let* $P = \bigcap_{i=0}^{n-1} \mathcal{H}_i$ *be a cyclic* n*-gon with sides* $\gamma_0, \ldots, \gamma_{n-1}$ *and center* v*, such that* $\gamma_i \subset \delta_i \doteq \partial\mathcal{H}_i$ *for each* i *and let* $\mathcal{H}'_i$ *be the half-space of* $\mathbb{H}^2$ *opposite* $\mathcal{H}_i$*. If* P *is not centered then there is a unique* $i_0 \in \{0, \ldots, n-1\}$ *such that* $v \in \mathcal{H}'_{i_0}$*. If* $T_{i_0} \subset \mathcal{H}'_{i_0}$ *is the triangle determined by* v *and* $\partial\gamma_{i_0}$ *then* $T_{i_0} \cap P = \gamma_{i_0}$*, and* $P \cup T_{i_0}$ *is a convex polygon.*

Proof. Let C be the circle containing the vertices of P, centered at v, and let B be the closed disk bounded by C. For each i let $C_i = C \cap \mathcal{H}'_i$, a closed subarc of C meeting γ_i at its endpoints, and let $B_i = \mathcal{H}'_i \cap B$, a convex subset of B bounded by $\gamma_i \cup C_i$ and homeomorphic to a disk. Then by construction, $B_i \cap P = \gamma_i$ for each i, $B_i \cap B_j$ is either empty or the single vertex $\gamma_i \cap \gamma_j$ for $j \neq i$, and $B = P \cup \bigcup_{i=0}^{n-1} B_i$.

Since v is not a vertex of P, it is contained in at most one of the B_i, and if P is non-centered then it is in some B_{i_0} (since in this case $v \notin int\, P$). This establishes the first claim above. If $v \in \gamma_{i_0}$ then the triangle T_{i_0} described above is degenerate: it equals γ_{i_0}, and hence $P \cup T_{i_0} = P$ is a convex polygon. Let us therefore assume below that $v \in B_{i_0} - \gamma_{i_0}$.

After re-numbering the γ_i if necessary, we may assume that for each i, γ_i intersects γ_{i+1} in a vertex x_i of P (taking $i+1$ modulo n). In particular, $\partial\gamma_{i_0} = x_{i_0-1} \cup x_{i_0}$. Let δ_- be the geodesic containing v and x_{i_0-1}, and let δ_+ be the geodesic containing v and x_{i_0}. Each of $\delta_\pm \cap B_{i_0}$ is a geodesic arc that contains v and hence intersects γ_{i_0} in a single point. Therefore $\delta_\pm \cap B_{i_0} = \delta_\pm \cap B$ is an arc with two endpoints in C_{i_0}.

It follows from the paragraph above that $P \cap \delta_\pm$ is a single point in ∂P, so $int\, P$ is contained in a complementary component to each of $\delta_\pm$. Let $\mathcal{H}_\pm$ be the closure of the component of $\mathbb{H}^2 - \delta_\pm$ containing $int\, P$. Then $P \subset \mathcal{H}_- \cap \mathcal{H}_+$. In particular, $x_{i_0} \subset \mathcal{H}_-$ and $x_{i_0-1} \subset \mathcal{H}_+$, so also $T_{i_0} \subset \mathcal{H}_- \cap \mathcal{H}_+$.

Let us finally note that $B_{i_0} \subset \mathcal{H}_j$ for each $j \neq i_0$, since $B_{i_0} \subset B$ and $B_j = \mathcal{H}'_j \cap B$ intersects B_{i_0} in at most a single point. This implies in particular that $T_{i_0} \subset \bigcap_{j \neq i_0} \mathcal{H}_j$, and hence that $P \cup T_{i_0} \subset \mathcal{H}_- \cap \mathcal{H}_+ \cap \left(\bigcap_{j \neq i_0} \mathcal{H}_j\right)$. Using frontiers as in the proof of Lemma 1.4, we find that equality holds, and hence by construction that $P \cup T_{i_0}$ is a convex polygon. □

Lemma 1.6. *Let* P *be a cyclic* n*-gon with center* v*, radius* J*, and sides* $\gamma_0, \ldots, \gamma_{n-1}$*. For* $i \in \{0, \ldots, n-1\}$*, let* T_i *be the triangle in* $\mathbb{H}^2$ *with vertex set* $\{v\} \cup \partial\gamma_i$*.*

- *If* P *is centered, then* $P = \bigcup_{i=0}^{n-1} T_i$*.*
- *If* P *is not centered, then* $P \cup T_{i_0} = \bigcup_{i \neq i_0} T_i$*, where* i_0 *is as in Lemma 1.5.*
- *If* P *is not centered but* $v \in P$*, then* $T_{i_0} = \gamma_{i_0} \subset P$ *and each conclusion above holds.*

In the first and last cases above, for $i \neq j$*,* $T_i \cap T_j = \{v\}$ *if* $\gamma_i \cap \gamma_j = \emptyset$ *and is the edge joining* $\gamma_i \cap \gamma_j$ *to* v *otherwise. This holds in the second case for distinct* i *and* j*, neither equal to* i_0*.*

Proof. Suppose first that P is centered. Since P is convex and $v \in P$, P contains the geodesic arc joining v to each endpoint of γ_i. Together with γ_i, these constitute the edges of T_i, and hence $T_i \subset P$. For any $x \in P - \{v\}$, let δ_x be the intersection with P of the geodesic ray from v in the direction of x. Then δ_x is an arc containing x with one endpoint at v and the other in ∂P, and therefore in γ_i for some i. Since T_i is convex, $\delta_x \subset T_i$ and hence $x \in T_i$. It follows that $P = \bigcup T_i$.

Since $v \in int\, P$, for x as above it follows that $\delta_x \cap \partial P$ is the endpoint $\delta_x \cap \gamma_i$. If x is also in T_j for some $j \neq i$, then $\delta_x \cap T_j$ is a sub-arc of δ_x containing x, with one endpoint at v and the other also in the frontier of T_j. If this endpoint is in γ_j, then since $\gamma_j \subset \partial P$ the first sentence implies that it is in $\gamma_j \cap \gamma_i$. But the endpoint of $\delta_x \cap T_j$ oposite v must be in γ_j, since the other two edges of T_j are arcs that join v to an endpoint of γ_j.

Now take P to be non-centered. With i_0 and T_{i_0} as in Lemma 1.5, since $P \cup T_{i_0}$ is convex and contains $\partial\gamma_i$ for each i, and also v, it contains T_i for each i. Since $T_{i_0} \cap P = \gamma_{i_0}$, the frontier of $T_{i_0} \cup P$ is the union of the γ_i, for $i \neq i_0$, with the sides of T_{i_0} containing v. If j_0 and j_1 are such that $\partial\gamma_{i_0} = (\gamma_{i_0} \cap \gamma_{j_0}) \cup (\gamma_{i_0} \cap \gamma_{j_1})$, then the sides of T_{i_0} containing v are contained in $T_{j_0} \cup T_{j_1}$, and hence $P \cup T_{i_0} = \bigcup_{i \neq i_0} T_i$.

In this case, T_{j_0} and T_{j_1} each intersects $\partial(P \cup T_{i_0})$ in a union of two edges. For $j \neq j_0$, $T_j \cap \partial(P \cup T_{i_0}) = \{v\} \cup \gamma_j$, and arguing as in the centered case we characterize $T_i \cap T_j$ for distinct i and j, neither equal to i_0. If P is not centered but $v \in P$, then $v \in \gamma_{i_0}$ by Lemma 1.5. Hence $T_{i_0} \subset \gamma_{i_0}$ by definition, and hence $P = P \cup T_{i_0} = \bigcup_{i \neq i_0} T_i = \bigcup T_i$. □

The functions below record the angles of some constituents of the decomposition above.

Lemma 1.7. *For $d > 0$ and $J \geq d/2$, define $A_d(J) \in (0, \pi]$ and $B_d(J) \in [0, \pi/2)$ as follows:*

$$A_d(J) = \cos^{-1}\left(1 - \frac{2\sinh^2(d/2)}{\sinh^2 J}\right) \qquad\qquad B_d(J) = \cos^{-1}\left(\frac{\tanh(d/2)}{\tanh J}\right)$$

The hyperbolic triangle with two sides of length J and one of length d has angle $A_d(J)$ at its vertex opposite the side with length d, and angle $B_d(J)$ at each other vertex.

To justify the statement above, we record the following well known fact: $0 < a \leq b \leq c$ with $c \leq a + b$ uniquely determines a triangle in $\mathbb{H}^2$ with sides of length a, b and c, up to isometries of $\mathbb{H}^2$. If $c = a + b$, then such a triangle is degenerate, contained in a single geodesic arc.

Proof. Let P be an isosceles triangle in $\mathbb{H}^2$ with two sides of length J and one of length d. The hyperbolic law of cosines (1.2.1) yields the equation below, relating J and d to the angle α of P at the vertex opposite the side γ with length d:

$$\cosh d = \cosh^2 J - \sinh^2 J \cos\alpha$$

Solving this equation for $\cos\alpha$ yields:

$$\cos\alpha = \frac{\cosh^2 J - \cosh d}{\sinh^2 J} = 1 - \frac{\cosh d - 1}{\sinh^2 J},$$

and since $\sinh(d/2) = \sqrt{(\cosh d - 1)/2}$ we find that $\alpha = A_d(J)$. If β is the angle of P at an endpoint of γ, the hyperbolic law of cosines gives:

$$\cosh J = \cosh J \cosh d - \sinh J \sinh d \cos\beta,$$

Solving for β and using hyperbolic trigonometric identities gives $\beta = B_d(J)$. □

Lemma 1.8. *For fixed* $d > 0$, $A_d(J)$ *strictly decreases on* $[d/2, \infty)$ *with* $A_d(d/2) = \pi$ *and* $\lim_{J\to\infty} A_d(J) = 0$. *For fixed* $J > 0$ *and* $0 < d < d' \leq 2J$, $A_d(J) < A_{d'}(J)$.

Proof. The function $1 - 2\sinh^2(d/2)/\sinh^2 J$ strictly increases in J for fixed d, and strictly decreases in d for fixed J. Thus since $\cos^{-1}$ is decreasing, $A_d(J)$ decreases in J for fixed d and increases in d for fixed J. Furthermore, $1 - 2\sinh^2(d/2)/\sinh^2 J$ takes a value of -1 at $J = d/2$ and limits to 0 as $J \to \infty$. Corresponding values for $A_d(J)$ again follow from properties of the inverse cosine. □

Below we will use the decomposition from Lemma 1.6 to produce equations relating the radius of a cyclic n-gon to its side lengths. In proving this it will be helpful to use the *unit tangent space* $UT_x\mathbb{H}^2 \doteq \{\eta \in T_x\mathbb{H}^2 \,|\, \langle\eta, \eta\rangle = 1\}$ at $x \in \mathbb{H}^2$. With the metric given by $d(\eta, \eta') = \cos^{-1}\langle\eta, \eta'\rangle$, $UT_x\mathbb{H}^2$ is isometric to the unit circle in the Euclidean plane.

Fact 1.9. If half-planes $\mathcal{H}$ and $\mathcal{H}'$ have frontier geodesics that meet at x, the set of vectors in $UT_x\mathbb{H}^2$ that point into $\mathcal{H} \cap \mathcal{H}'$ is isometric to an interval embedded in $UT_x\mathbb{H}^2$, with length equal to the interior angle at x between $\mathcal{H}$ and $\mathcal{H}'$ (see Definition 1.2).

If x is a vertex of a polygon P, let us denote by $UT_xP \subset UT_x\mathbb{H}^2$ the interval determined by $\mathcal{H}$ and $\mathcal{H}'$ as above, where $\mathcal{H}$ and $\mathcal{H}'$ are the half-planes containing P and bounded by geodesics containing the edges of P that intersect at x. Fact 1.9 and Definition 1.2 imply that UT_xP has length $\angle_x P$.

Lemma 1.10. *Let* P *be a cyclic* n*-gon in* $\mathbb{H}^2$ *with center* v, *radius* J, *and cyclically ordered vertices* $x_0, \ldots, x_{n-1}$, *and sides* γ_i *with* $\partial\gamma_i = x_{i-1} \cup x_i$ *and* $\ell(\gamma_i) = d_i$ *for each* i.

- *If* P *is centered then* $\sum_{i=0}^{n-1} A_{d_i}(J) = 2\pi$ *and for each* i, $\angle_{x_i} P = B_{d_i}(J) + B_{d_{i+1}}(J)$.
- *If not,* $A_{d_{i_0}}(J) = \sum_{i\neq i_0} A_{d_i}(J)$ *with* i_0 *as in Lemma 1.5. For* $i \notin \{i_0, i_0 - 1\}$, $\angle_{x_i} P = B_{d_i}(J) + B_{d_{i+1}}(J)$; *but* $\angle_{x_{i_0}} P = B_{d_{i_0+i}}(J) - B_{d_{i_0}}(J)$ *and* $\angle_{x_{i_0-1}} P = B_{d_{i_0-1}}(J) - B_{d_{i_0}}(J)$.
- *If* P *is not centered but* $v \in P$ *then* $A_{d_{i_0}}(J) = \pi$, $B_{d_{i_0}}(J) = 0$, *and each assertion above holds.*

Above $A_{d_i}(J)$ *and* $B_{d_i}(J)$ *are as in Lemma 1.7, and we take* $i \pm 1$ *modulo* n *when appropriate.*

Proof. For P as above, let $\{T_i\}_{i=0}^{n-1}$ be the decomposition from Lemma 1.6. Lemma 1.7 implies that $A_{d_i}(J) = \angle_v T_i$, and $B_{d_i}(J) = \angle_{x_i} T_i = \angle_{x_{i-1}} T_i$, for each i.

If P is centered, then since $v \in int\, P$ there is an open disk U centered at v with small enough radius that $U \subset P$. Each $\eta \in UT_v\mathbb{H}^2$ determines a geodesic ray from v with an initial interval in U, and hence in T_i for some i, by Lemma 1.6. Therefore $\eta \in UT_vT_i$, and it follows that $UT_v\mathbb{H}^2 \subset \bigcup_{i=0}^{n-1} UT_vT_i$. Lemma 1.6 also implies that for $i \neq j$, T_i intersects T_j in at most an edge containing v; hence UT_vT_i intersects UT_vT_j in at most a single endpoint. Since $UT_v\mathbb{H}^2$ is the Euclidean unit circle its circumference is 2π. This is the sum of the lengths of the UT_vT_i, so the observation above the lemma gives $2\pi = \sum_{i=0}^{n-1} \angle_v T_i = \sum_{i=0}^{n-1} A_{d_i}(J) = 2\pi$.

We will use the following fact repeatedly: if a vertex x of a convex polygon Q has a neighborhood U such that $Q \cap U = \bigcup_{i=0}^{n-1} P_i \cap U$, where the P_i are non-overlapping convex polygons containing x, then $\angle_x Q = \sum_{i=0}^{n-1} \angle_x P_i$. This is proved as in the paragraph above, by observing that the $UT_x P_i$ are non-overlapping intervals in $UT_x \mathbb{H}^2$ whose union is $UT_x Q$.

For $i \in \{0, \dots, n-1\}$, by Lemma 1.6 x_i is contained in T_i, T_{i+1}, and no other T_j. Thus there is a disk U_i centered at x_i small enough that $U_i \cap P = (U_i \cap T_i) \cup (U_i \cap T_{i+1})$, so by the fact above $\angle_{x_i} P = \angle_{x_i} T_i + \angle_{x_i} T_{i+1} = B_{d_i}(J) + B_{d_{i+1}}(J)$.

If P is non-centered and $v \notin P$, then v has an open neighborhood U in $\mathbb{H}^2 - P$, and it follows from Lemma 1.6 that $T_{i_0} \cap U = \bigcup_{i \neq i_0} T_i \cap U$. The fact above thus implies:

$$A_{d_{i_0}}(J) = \angle_v T_{i_0} = \sum_{i \neq i_0} \angle_v T_i = \sum_{i \neq i_0} A_{d_i}(J)$$

In this case, the vertex x_{i_0} of P is contained in T_{i_0}, T_{i_0+1} and no others by Lemma 1.6. Thus if U is an open neighborhood of x_{i_0} small enough to miss $\bigcup_{i \neq i_0, i_0+1} T_i$, that lemma implies that $U \cap (P \cup T_{i_0}) = U \cap T_{i_0-1}$. The fact above thus gives $\angle_{x_{i_0}} P + \angle_{x_{i_0}} T_{i_0} = \angle_{x_{i_0}} T_{i_0+1}$, giving $\angle_{x_{i_0}} P = B_{d_{i_0+1}}(J) - B_{d_{i_0}}(J)$. A similar argument gives a similar conclusion at x_{i_0-1}, and arguing as in the centered case gives $\angle_{x_i} P = B_{d_i}(J) + B_{d_{i-1}}(J)$ for $i \notin \{i_0, i_0 - 1\}$.

If P is non-centered but $v \in P$, then $v \in \gamma_{i_0}$ by Lemma 1.5. Since v is in particular equidistant from the endpoints of γ_{i_0}, $d_{i_0} = 2J$, so Lemma 1.8 implies that $A_{d_{i_0}}(J) = \pi$. From the definition in Lemma 1.7, $B_{d_{i_0}}(J) = 0$. Note that this agrees with the geometric assertion from Lemma 1.6, that $T_{i_0} = \gamma_{i_0} \subset P$.

In this case, there is a neighborhood U of v small enough that $U \cap P = U \cap \mathcal{H}_{i_0}$, where $\mathcal{H}_{i_0}$ is the half-space containing P and bounded by the geodesic containing γ_{i_0}. Therefore $UT_{x_{i_0}}(P \cup T_{i_0}) = UT_{x_{i_0}} \mathcal{H}_{i_0}$ is a segment in $UT_{x_{i_0}} \mathbb{H}^2$ with length π. (This is a feature of half-spaces, as can be easily verified.) Thus Lemma 1.6 and the fact above imply in this case that $\sum_{i \neq i_0} A_{d_i}(J) = \pi = A_{d_{i_0}}(J)$, and hence also $\sum_{i=0}^{n-1} A_{d_i}(J) = 2\pi$.

To establish the values of $\angle_{x_i} P$ in this case we argue as in the case $v \notin P$ and note that $B_{d_{i_0+1}}(J) - B_{d_{i_0}}(J) = B_{d_{i_0+1}}(J) + B_{d_{i_0}}(J)$ since $B_{d_{i_0}}(J) = 0$. □

Corollary 1.11. *For $n \geq 3$, if P is a non-centered cyclic n-gon with sides $\gamma_0, \dots, \gamma_{n-1}$ then for i_0 as identified in Lemma 1.5, γ_{i_0} is the unique longest side of P.*

Proof. If J is the radius, then since $A_{d_{i_0}}(J) = \sum_{i \neq i_0} A_{d_i}(J)$ and all summands are positive, $A_{d_{i_0}}(J) > A_{d_i}(J)$ for each $i \neq i_0$. By Lemma 1.8, this implies that $d_{i_0} > d_i$ for each $i \neq i_0$. □

2. Existence and uniqueness of cyclic and horocyclic polygons

The main result of this section is Proposition 2.7, which asserts existence and uniqueness of a cyclic n-gon P with side length collection $(d_0, \dots, d_{n-1}) \in (\mathbb{R}^+)^n$ if and only if $\sinh(d_i/2) < \sum_{j \neq i} \sinh(d_j/2)$ for each i. An additional inequality determines whether P is centered: assuming that d_0 is maximal among the d_i, P is centered if and only if $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) > \pi$. Analogously, by Proposition 2.9 there is a (unique) horocyclic n-gon with the above side length collection if and only if $\sinh(d_i/2) = \sum_{j \neq i} \sinh(d_j/2)$ for some i.

The most important step in proving both existence and uniqueness of cyclic polygons is to show that a side length collection satisfying the first or both inequalities above determines a unique radius J satisfying the appropriate equation from Lemma 1.10. This is the content of Lemma 2.2, proved by carefully examining a certain ratio of derivatives of the functions $A_d(J)$ from Lemma 1.7. With the radius thus determined, P can be reconstituted using the decomposition from Lemma 1.6. Below we record some key observations about the $A_d(J)$.

Lemma 2.1. *The map $(d, J) \mapsto A_d(J)$ is differentiable on $U = \{(d, J) \mid d > 0, J > d/2\}$ and continuous on $\overline{U}$, with $\lim_{J\to d_0/2^+} A'_{d_0}(J) = -\infty$. For $n \geq 3$ and $\mathcal{D} = \{d_0, \ldots, d_{n-1}\} \subset \mathbb{R}^+$ such that $d_0 \geq d_i$ for each $i > 0$:*

$$\lim_{J\to\infty} \frac{\sum_{i=1}^{n-1} A_{d_i}(J)}{A_{d_0}(J)} = \lim_{J\to\infty} \frac{\sum_{i=1}^{n-1} A'_{d_i}(J)}{A'_{d_0}(J)} = \frac{\sum_{i=1}^{n-1} \sinh(d_i/2)}{\sinh(d_0/2)}$$

Furthermore, the derivative ratio $\left(\sum_{i=1}^{n-1} A'_{d_i}(J)\right) / A'_{d_0}(J)$ is strictly increasing on $(d_0/2, \infty)$.

Proof. That $(d, J) \mapsto A_d(J)$ is differentiable on U and continuous on $\overline{U}$ follows from elementary properties of the inverse cosine and hyperbolic sine. Below we fix d and record the derivative of $A_d(J)$ with respect to J:

$$A'_d(J) = -2\frac{\cosh J}{\sinh J}\sqrt{\frac{\cosh d - 1}{2\sinh^2 J - \cosh d + 1}} = -2\frac{\cosh J}{\sinh J}\frac{\sinh(d/2)}{\sqrt{\sinh^2 J - \sinh^2(d/2)}} \tag{2.1.1}$$

Inspecting this formula shows that $A'_d(J)$ approaches $-\infty$ as J approaches $d/2$ from the right. Substituting the d_i and taking the ratio under consideration yields:

$$\frac{\sum_{i=1}^{n-1} A'_{d_i}(J)}{A'_{d_0}(J)} = \sum_{i=1}^{n-1} \frac{\sinh(d_i/2)}{\sinh(d_0/2)}\sqrt{\frac{\sinh^2 J - \sinh^2(d_0/2)}{\sinh^2 J - \sinh^2(d_i/2)}} \tag{2.1.2}$$

Since $A_{d_i}(J) \to 0$ as $J \to \infty$ for each i, L'Hopital's rule implies that

$$\lim_{J\to\infty} \frac{\sum_{i=1}^{n-1} A_{d_i}(J)}{A_{d_0}(J)} = \lim_{J\to\infty} \frac{\sum_{i=1}^{n-1} A'_{d_i}(J)}{A'_{d_0}(J)}$$

The claimed description of this limit follows upon inspecting (2.1.2). A slight rewrite yields:

$$\frac{\sum_{i=1}^{n-1} A'_{d_i}(J)}{A'_{d_0}(J)} = \sum_{i=1}^{n-1} \frac{\sinh(d_i/2)}{\sinh(d_0/2)}\sqrt{1 - \frac{\sinh^2(d_0/2) - \sinh^2(d_i/2)}{\sinh^2 J - \sinh^2(d_i/2)}} \tag{2.1.3}$$

From this it is clear that the ratio of derivatives strictly increases with J. □

Lemma 2.2. *For $n \geq 3$, suppose for $\mathcal{D} = \{d_0, \ldots, d_{n-1}\} \subset \mathbb{R}^+$ that for each i, $\sinh(d_i/2) < \sum_{j\neq i} \sinh(d_j/2)$. If d_{i_0} is the largest element of $\mathcal{D}$, then:*

1. *If $\sum_{i\neq i_0} A_{d_i}(d_{i_0}/2) \geq \pi$, there exists $J_0 \geq d_{i_0}/2$ such that $\sum_{i=0}^{n-1} A_{d_i}(J_0) = 2\pi$; or*
2. *If $\sum_{i\neq i_0} A_{d_i}(d_{i_0}/2) \leq \pi$, there exists $J_1 \geq d_{i_0}/2$ such that $A_{d_{i_0}}(J_1) = \sum_{i\neq i_0} A_{d_i}(J_1)$, and $A_{d_{i_0}}(J) < \sum_{i\neq i_0} A_{d_i}(J)$ for each $J > J_1$.*

The solution in either case above is unique. In case (1) there is no J_1 satisfying the conclusion of case (2), and vice-versa, unless $\sum_{i\neq i_0} A_{d_i}(d_{i_0}/2) = \pi$. In this case $J_0 = J_1 = d_{i_0}/2$.

Proof. Re-numbering if necessary, we will assume $i_0 = 0$. If $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) \geq \pi$ then since $A_{d_0}(d_0/2) = \pi$, we have $\sum_{i=0}^{n-1} A_{d_i}(d_0/2) \geq 2\pi$. By Lemma 1.8, $\sum_{i=0}^{n-1} A_{d_i}(J)$ strictly decreases in J with a horizontal asymptote of 0, so in this case there is a unique $J_0 \geq d_0/2$ such that $\sum_{i=0}^{n-1} A_{d_i}(J_0) = 2\pi$. Note that if $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) = \pi$ then $J_0 = d_0/2$, and otherwise $J_0 > d_0/2$.

If $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) = \pi$ then $J_1 = J_0 = d_0/2$ also satisfies $A_{d_0}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1)$. If $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) < \pi$, then the ratio of functions $\left(\sum_{i=1}^{n-1} A_{d_i}(J)\right)/A_{d_0}(J)$ takes a value less than 1 at $J = d_0/2$, but by Lemma 2.1 its limit is greater than 1 as $J \to \infty$. It follows that in this case there exists $J_1 > d_0/2$ such that $A_{d_0}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1)$.

Note that $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) \leq \pi$ implies that $d_i < d_0$ for all $i > 0$, since $A_{d_0}(d_0/2) = \pi$ and $d \mapsto A_d(J)$ is increasing for fixed J. Lemma 2.1 thus implies in this case that the derivative ratio $\left(\sum_{i=1}^{n-1} A'_{d_i}(J)\right)/A'_{d_0}(J)$ is less than one near $J = d_0/2$ and limits to a value greater than 1 as $J \to \infty$. Since it is strictly increasing, there exists $J_2 > d_0/2$ such that A_{d_0} decreases faster than $\sum_{i=1}^{n-1} A_{d_i}$ on $(d_0/2, J_2)$ and more slowly on (J_2, ∞).

Assuming that $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) \leq \pi$, let J_1 be the smallest solution to $A_{d_0}(J) = \sum_{i=1}^{n-1} A_{d_i}(J)$. If $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) = \pi$ then $J_1 = d_0/2$, and if $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) < \pi$ then $A_{d_0} > \sum_{i=1}^{n-1} A_{d_i}$ on $[d_0/2, J_1)$. Since $\sum_{i=1}^{n-1} A_{d_i}$ decreases faster than A_{d_0} on (J_2, ∞), we must have $J_1 \leq J_2$. In fact, if $A_{d_0}(J)$ were equal to $\sum_{i=1}^{n-2} A_{d_i}(J)$ for any $J \geq J/2$ then A_{d_0} and $\sum_{i=1}^{n-1} A_{d_i}$ would have different limits as $J \to \infty$, a contradiction. In particular, $J_1 < J_2$. But since A_{d_0} decreases faster than $\sum_{i=1}^{n-1} A_{d_i}$ on (J_1, J_2), J_1 is the unique solution to $A_{d_0}(J) = \sum_{i=1}^{n-1} A_{d_i}(J)$ on this interval, and it follows that J_1 is the only solution on $[d_0/2, \infty)$.

We now address the mutual-exclusivity property that is the final claim of the lemma. If $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) < \pi$ then $\sum_{i=0}^{n-1} A_{d_i}(d_0/2) < 2\pi$, so since $\sum_{i=0}^{n-1} A_{d_i}(J)$ decreases in J there does not exist J_0 satisfying $\sum_{i=0}^{n-1} A_{d_i}(J_0) = 2\pi$. Thus in case (2) above the conclusion of case (1) does not also hold unless $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) = \pi$. We claim that in case (1) there does not exist J_1 such that $A_{d_0}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1)$, unless $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) = \pi$.

Assume that $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) > \pi$. Again, the derivative ratio $\left(\sum_{i=1}^{n-1} A'_{d_i}(J)\right)/A'_{d_0}(J)$ takes values greater than one on some interval (J_2, ∞). Taking J_2 to be minimal with this property, we may in this case have $J_2 = d_0/2$, since values of $\left(\sum_{i=1}^{n-1} A'_{d_i}(J)\right)/A'_{d_0}(J)$ near $d_0/2$ are not clear. However, if $J_2 > d_0/2$ then since the derivative ratio is strictly increasing, its values are less than 1 on $(d_0/2, J_2)$.

There is no $J_1 \in [J_2, \infty)$ such that $A_{d_0}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1)$ since this would contradict the fact that A_{d_0} and $\sum_{i=1}^{n-1} A_{d_i}$ have the same limits as $J \to \infty$ but $\sum_{i=1}^{n-1} A_{d_i}$ decreases faster on (J_1, ∞). On the other hand, there is no $J_1 \in [d_0/2, J_2)$ satisfying $A_{d_0}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1)$, since $A_{d_0}(d_0/2) = \pi < \sum_{i=1}^{n-1} A_{d_i}(d_0/2)$ and A_{d_0} decreases faster on $(d_0/2, J_2)$. It follows that in this case there is no solution to $A_{d_0}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1)$. □

Lemma 2.3. *For $n \geq 3$, let $\mathcal{D} = \{d_0, \ldots, d_{n-1}\} \subset \mathbb{R}^+$, and suppose $d_0 \geq d_i$ for each $i > 0$. If $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) \geq \pi$, then $\sinh(d_0/2) < \sum_{i=1}^{n-1} \sinh(d_i/2)$.*

Proof. Rearranging the definition of $A_d(J)$ (cf. Lemma 1.7) yields the following identity:

$$\sinh^2(d/2) = \sinh^2 J \cdot \frac{1-\cos A_d(J)}{2} \qquad \Rightarrow \qquad \sinh(d/2) = \sinh J \sin(A_d(J)/2)$$

Inserting d_i for d and $d_0/2$ for J above, for $1 \leq i \leq n-1$, and summing the results yields:

$$\sum_{i=1}^{n-1} \sinh(d_i/2) = \sinh(d_0/2) \cdot \left(\sum_{i=1}^{n-1} \sin(A_{d_i}(d_0/2)/2) \right)$$

To finish the proof we record the following simple fact.

Claim 2.3.1. If $0 < \theta_i \leq \pi/2$ for $1 \leq i \leq k$ $(k \geq 2)$ and $\sum_{i=1}^k \theta_i \geq \pi/2$, then $\sum_{i=1}^k \sin\theta_i > 1$.

Proof. This follows immediately if $\theta_i = \pi/2$ for any i, so we will assume henceforth that $0 < \theta_i < \pi/2$ for all i. Let us first suppose that $\alpha + \beta \geq \pi/2$ for $0 < \alpha, \beta < \pi/2$. Then since $\beta \geq \pi/2 - \alpha$ it follows that $\sin\beta \geq \sin(\pi/2 - \alpha) = \cos\alpha$. Thus:

$$\sin\alpha + \sin\beta \geq \sin\alpha + \cos\alpha > \sin^2\alpha + \cos^2\alpha = 1$$

Using the above, we conclude that $\sin\theta_1 + \sin(\theta_2 + \dots \theta_k) > 1$. We will complete the proof by inductively applying the following observation: for $0 < \alpha, \beta < \pi/2$,

$$\sin(\alpha + \beta) = \sin\alpha\cos\beta + \cos\alpha\sin\beta < \sin\alpha + \sin\beta$$

Therefore $\sin\theta_2 + \ldots + \sin\theta_k > \sin(\theta_2 + \ldots + \theta_k)$, and the claim is proved. □

We note that for $i > 0$, since $d_0 \geq d_i$ and A_{d_i} is decreasing, $A_{d_i}(d_0/2) \leq A_{d_i}(d_i/2) = \pi$. Thus if $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) \geq \pi$, then using the claim and the equation above it we find that $\sum_{i=1}^{n-1} \sinh(d_i/2) > \sinh(d_0/2)$. □

Henceforth we let S_n be the symmetric group on n-letters, acting on $\{0, 1, \ldots, n-1\}$.

Lemma 2.4. *For $n \geq 3$, $\mathcal{D} = \{d_0, \ldots, d_{n-1}\} \subset \mathbb{R}^+$, with largest element d_0, is the collection of side lengths of a cyclic n-gon in $\mathbb{H}^2$ if and only if $\sinh(d_0/2) < \sum_{i=1}^{n-1} \sinh(d_i/2)$. If so then for any $\sigma \in S_n$ there is a cyclic n-gon P with side length collection $\mathcal{D}$ and a point $x \in \partial P$ so that starting from x and moving along ∂P in one direction, the edges encountered have lengths $d_{\sigma(0)}, d_{\sigma(1)}, \ldots, d_{\sigma(n)}$ in that order. Any such P is centered if and only if $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) > \pi$; or equivalently, if and only if $\sum_{i=0}^{n-1} A_{d_i}(d_0/2) > 2\pi$.*

Proof. We will prove the lemma by reverse-engineering the decomposition from Lemma 1.6. First suppose that $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) > \pi$, so by Lemma 2.2 there exists J_0 with $\sum_{i=0}^{n-1} A_{d_i}(J_0) = 2\pi$. For each i, let T_i be an isosceles triangle with two sides of length J_0 and one of length d_i. Then by Lemma 1.7, $A_{d_i}(J_0)$ is the angle of T_i at the vertex v_i opposite the side with length d_i. We fix a choice of $\sigma \in S_n$ and use it to determine the following inductive construction.

Let $v = v_{\sigma(0)}$, let $f_0 \colon \mathbb{H}^2 \to \mathbb{H}^2$ be the identity map, and choose an isometry f_1 of $\mathbb{H}^2$ so that $f_1(v_{\sigma(1)}) = v$ and $f_1(T_{\sigma(1)}) \cap T_{\sigma(0)}$ is an edge of each (with length J_0, since it contains v). For $1 < i \leq n-1$, assume that an isometry f_{i-1} has been chosen so that $f_{i-1}(v_{\sigma(i-1)}) = v$ and $f_{i-1}(T_{\sigma_{i-1}}) \cap f_{i-2}(T_{\sigma_{i-2}})$ is an edge of each, and let f_i take $v_{\sigma(i)}$ to v so that $f_i(T_{\sigma(i)}) \cap$

$f_{i-1}(T_{\sigma(i-1)})$ is an edge of each that is not contained in $f_{i-2}(T_{\sigma(i-2)})$. Induction yields a collection of triangles $f_i(T_{\sigma(i)})$, $i \in \{0, \ldots, n-1\}$ all sharing the vertex v.

For $i < n-1$ let x_i be the endpoint opposite v on the edge $f_i(T_{\sigma(i)}) \cap f_{i+1}(T_{\sigma(i+1)})$. Since these edges all have length J_0, all of the x_i, $i \in \{0, \ldots, n-2\}$, lie on a circle of radius J_0 centered at v. Furthermore, $d_{\sigma(i)} = d(x_{i-1}, x_i)$ for $1 \leq i < n-1$ by construction. We claim that $f_{n-1}(T_{\sigma(n-1)})$ intersects $f_0(T_{\sigma(0)})$ in an edge containing v, with other endpoint x_{n-1} neither equal to x_0 nor x_{n-2}.

Recall from Fact 1.9 and the observation below it that for each i, $UT_v f_i(T_{\sigma(i)})$ is an interval of length $A_{d_{\sigma(i)}}(J)$ isometrically embedded in $UT_v\mathbb{H}^2$. Since $T_{\sigma(0)} = f_0(T_{\sigma(0)})$ intersects $f_1(T_{\sigma(1)})$ in the edge containing x_0, $UT_v T_{\sigma(0)}$ intersects $UT_v f_1(T_{\sigma(1)})$ in an endpoint of each: the unit tangent vector η_0 to this edge at v, pointing in the direction of x_0. For $1 < i \leq n-1$, $UT_v f_i(T_{\sigma(i)})$ similarly intersects $UT_v f_{i-1}(T_{\sigma(i-1)})$ in the endpoint η_i opposite $\eta_{i-1} = UT_v f_{i-1}(T_{\sigma(i-1)}) \cap UT_v f_{i-2}(T_{\sigma(i-2)})$.

Since $\sum_{i=0}^{n-1} A_{d_{\sigma(i)}}(J) = 2\pi$, a locally isometric immersion of $[0, 2\pi]$ into $UT_v\mathbb{H}^2$ is determined by restricting on $[0, A_{d_{\sigma(0)}}(J)]$ to the embedding of $UT_v T_{\sigma(0)}$ and to the embedding of $UT_v f_i(T_{\sigma(i)})$ on $[\sum_{j=0}^{i-1} A_{d_{\sigma(j)}}(J), \sum_{j=0}^{i} A_{d_{\sigma(j)}}(J)]$ for $0 < i \leq n-1$. Since $UT_v\mathbb{H}^2$ is the Euclidean unit circle, in particular with circumference 2π, this immersion is embedding on $(0, 2\pi)$ and sends 0 to the same point as 2π. Thus $UT_v T_{\sigma(0)}$ shares an endpoint η_{n-1} with $UT_v f_{n-1}(T_{\sigma(n-1)})$, neither equal to η_0 nor to η_{n-2}, and the claim follows.

Thus $\{x_0, x_1, \ldots, x_{n-1}\}$ is cyclically ordered on the circle of radius J_0 centered at v, with $d_{\sigma(i)} = d(x_{i-1}, x_i)$ for each i, taking $i-1$ modulo n. Applying Lemma 1.4 produces a cyclic polygon P with vertex set $\{x_i\}$ and side length collection $\mathcal{D}$, so that moving from $x \doteq x_{n-1}$ in the direction of x_0 on ∂P the edges encountered have lengths $d_{\sigma(0)}, \ldots, d_{\sigma(n-1)}$.

By the above, $\mathcal{D}$ is the side length collection of a centered polygon if $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) > \pi$; or equivalently, if $\sum_{i=0}^{n-1} A_{d_i}(d_0/2) > 2\pi$ (since $A_{d_0}(d_0/2) = \pi$, cf. Lemma 1.8). On the other hand, if P is a centered n-gon with radius J_0 and side length collection $\mathcal{D}$, then Lemma 1.10 implies that $\sum_{i=1}^{n-1} A_{d_i}(J_0) = 2\pi$ and $J_0 > d_0/2$, so $\sum_{i=0}^{n-1} A_{d_i}(d_0/2) > 2\pi$ since $A_d(J)$ decreases in J. Therefore $\mathcal{D}$ is the side length collection of a centered polygon P if and only if $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) > \pi$. In this case, Lemma 2.3 implies that $\sinh(d_0/2) < \sum_{i=1}^{n-1} \sinh(d_i/2)$, and Lemma 2.2 implies that there is no solution J_1 to the equation $A_{d_0}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1)$. Therefore by Lemma 1.10, every cyclic n-gon with side length collection $\mathcal{D}$ is centered.

Let us now suppose that $\sinh(d_0/2) = \sum_{i=1}^{n-1} \sinh(d_i/2)$ but that $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) \leq \pi$. Then Lemma 2.2 supplies J_1 such that $A_{d_0}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1)$. Again let $T_i \subset \mathbb{H}^2$ be an isosceles triangle with two sides of length J_1 and one of length d_i, so $A_{d_i}(J_1)$ is the angle of T_i at the vertex v_i opposite the side with length d_i. In this case, given $\sigma \in S_n$ let $i_0 \in \{0, \ldots, n-1\}$ satisfy $\sigma(i_0) = 0$. We again apply the inductive construction from the centered case, but this time to the collection $\{T_{\sigma(i_0+i)}\}_{i=1}^{n-1}$ (taking $i_0 + i$ modulo n), with base case $i = 1$.

This construction produces isometries $f_1 = id_{\mathbb{H}^2}, f_2, \ldots, f_{n-1}$ such that the $f_i(T_{\sigma(i_0+i)})$ share the vertex $v \doteq v_{\sigma(i_0+1)}$ for all i, and $f_i(T_{\sigma(i_0+i)})$ intersects $f_{i-1}(T_{\sigma(i_0+i-1)})$ in one edge containing v and $f_{i+1}(T_{\sigma(i_0+i+1)})$ in the other for $1 < i < n-1$.

For $1 \leq i < n-1$, let x_{i_0+i} be the endpoint opposite v on $f_i(T_{\sigma(i_0+i)}) \cap f_i(T_{\sigma(i_0+i+1)})$. Let x_{i_0} be the vertex of $T_{\sigma(i_0+1)}$ at distance J_1 from v but not equal to x_1, and let x_{i_0-1} be the vertex of $f_{n-1}(T_{\sigma(i_0+n-1)})$ at distance J_1 from v but not equal to x_{i_0+n-2}. By construction the x_i all lie on the circle of radius J_1 centered at v, and $d(x_{i_0+i-1}, x_{i_0+i}) = d_{\sigma(i_0+i)}$ for each $i > 0$ (taking indices modulo n above). It remains to determine $d(x_{i_0-1}, x_{i_0})$.

As in the centered case, there is a locally isometric immersion from a Euclidean interval $[0, A]$ to $UT_v\mathbb{H}^2$ that restricts to the isometric embedding of $UT_vT_{\sigma(i_0+1)}(J_1)$ on $[0, A_{d_{\sigma(i_0+1)}}(J_1)]$ and to $UT_vf_i(T_{\sigma(i)})$ on $[\sum_{j=1}^{i-1} A_{d_{\sigma(i_0+j)}}(J_1), \sum_{j=1}^{i} A_{d_{\sigma(i_0+j)}}(J_1)]$ for $1 < i \leq n-1$. Recalling that $\sigma(i_0) = 0$, we find that the length A of this interval satisfies:

$$A = \sum_{i=1}^{n-1} A_{d_{\sigma(i_0+i)}}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1) = A_{d_0}(J_1)$$

Since $A_{d_0}(J_1) < \pi$, in this case $[0, A]$ is embedded in $UT_v\mathbb{H}^2$. By construction the endpoints of the image are vectors η_{i_0} and η_{i_0-1} that point in the direction of x_{i_0} and x_{i_0-1}, respectively. If $\mathcal{H}_{i_0}$ is the half-space containing $T_{\sigma(i_0+1)}$ and bounded by the geodesic containing v and x_{i_0}, and $\mathcal{H}_{i_0-1}$ contains $f_{n-1}(T_{\sigma(i_0-1)})$ with boundary containing v and x_{i_0-1}, then by Fact 1.9 and the above the interior angle at v between $\mathcal{H}_{i_0}$ and $\mathcal{H}_{i_0-1}$ is $A_{d_0}(J_1)$. Applying the hyperbolic law of cosines to the triangle T_0 determined by v, x_{i_0}, and x_{i_0-1} gives $d(x_{i_0}, x_{i_0-1}) = d_0$.

Lemma 1.4 implies that $\{x_0, \ldots, x_{n-1}\}$ is the vertex collection of a cyclic polygon P with side length collection $\mathcal{D}$, so that moving from $x \doteq x_{n-1}$ in the direction of x_0 the edge lengths encountered are $d_{\sigma(0)}, \ldots, d_{\sigma(n-1)}$. It remains to show that if P is a cyclic n-gon with side length collection $\mathcal{D}$, then $\sinh(d_0/2) < \sum_{i=1}^{n-1} \sinh(d_i/2)$. We have already covered the case that $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) > \pi$, so let us assume below that $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) \leq \pi$. This implies in particular that $d_i < d_0$ for each $i > 0$, since $A_{d_0}(d_0/2) = \pi$.

Since $\lim_{J \to d_0/2^+} A'_{d_0}(J) = -\infty$ by Lemma 2.1, there is an open interval $(d_0/2, J_2)$ on which $|\sum_{i=1}^{n-1} A'_{d_i}(J)| < |A'_{d_0}(J)|$. If the maximal such J_2 is finite then $\sum_{i=1}^{n-1} \sinh(d_i/2) > \sinh(d_0/2)$, since $\left(\sum_{i=1}^{n-1} A'_{d_i}(J)\right) / A'_{d_0}(J)$ is strictly increasing and limits to $\left(\sum_{i=1}^{n-1} \sinh(d_i/2)\right) / \sinh(d_0/2)$ as $J \to \infty$. Let us therefore assume below that $J_2 = \infty$; that is, that A_{d_0} decreases faster than $\sum_{i=1}^{n-1} A_{d_i}$ on all of $[d_0/2, \infty)$.

If J_1 is the radius of P, then Lemma 1.10 implies that $A_{d_0}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1)$. Since $A'_{d_0} < \sum_{i=1}^{n-1} A'_{d_i}$ on $[J_1, \infty)$, it follows that

$$\lim_{J \to \infty} \sum_{i=1}^{n-1} A_{d_i}(J) - A_{d_0}(J) = \lim_{J \to \infty} \int_{J_1}^{J} \sum_{i=1}^{n-1} A'_{d_i} - A'_{d_0} > 0.$$

This is impossible, however, as each of A_{d_0} and $\sum_{i=1}^{n-1} A_{d_i}$ limits to 0 as $J \to \infty$. It follows that J_2 is finite, and hence that $\sinh(d_0/2) < \sum_{i=1}^{n-1} \sinh(d_i/2)$. □

Definition 2.5. The collection $\gamma_0, \ldots, \gamma_{n-1}$ of edges of a convex n-gon P in $\mathbb{H}^2$ is *cyclically ordered* if γ_i shares an endpoint with γ_{i+1} for each i taking $i+1$ modulo n. If so, taking $d_i = \ell(\gamma_i)$ for each i we say $(d_0, \ldots, d_{n-1}) \in (\mathbb{R}^+)^n$ is the corresponding *cyclically ordered side* *length* collection. (If P is cyclic, then $\{\gamma_i\}$ is cyclically ordered if and only if the collection $\{x_i \doteq \gamma_i \cap \gamma_{i+1}\}$ is cyclically ordered in the sense of Definition 1.3.)

If the collection $\{\gamma_i\}$ is cyclically ordered then it remains so if all indices are changed by a power of the cyclic permutation $\sigma\colon i \mapsto i+1 \,(\mathrm{mod}\, n)$, or by the permutation $\tau\colon i \mapsto n-i \,(\mathrm{mod}\, n)$. If there is some $k \in \{1,\dots,n-1\}$ such that $d'_i = d_{\sigma^k(i)}$ for all i we say the tuple $(d'_0,\dots,d'_{n-1})$ is obtained by *cyclically re-ordering* $(d_0,\dots,d_{n-1})$. Cyclic re-ordering is equivalent to starting one's edge count from a different point, and re-indexing by τ to counting in a different direction along ∂P.

If P has cyclically ordered side collection $\{\gamma_0,\dots,\gamma_{n-1}\}$, there is an orientation on ∂P such that γ_i points from its intersection with γ_{i-1} to γ_{i+1} for each i. Cyclically re-ordering preserves this orientation, whereas re-ordering by τ reverses it. Thus in the group $\langle\sigma,\tau\rangle < S_n$, isomorphic to the dihedral group D_{2n} of order $2n$, $\langle\sigma\rangle$ is the orientation-preserving subgroup. We will refer to this group by $\mathbb{Z}_n \triangleleft D_{2n}$.

Definition 2.6. For a convex n-gon $P \subset \mathbb{H}^2$, the *standard* orientation on ∂P is the boundary orientation that P gives it, when P inherits the standard orientation on $\mathbb{H}^2$. We will say that a cyclic order $\gamma_0,\dots,\gamma_{n-1}$ for the collection of sides of P (see Definition 2.5) is *standard* if for each i, γ_i points toward γ_{i+1} in the standard orientation on ∂P.

It is useful to note that for any fixed $i \in \{0,\dots,n-1\}$ the tuple $(d_0,\dots,d_{n-1})$ may be cyclically re-ordered to move d_i to the first position.

Proposition 2.7. *There is a cyclic n-gon in $\mathbb{H}^2$ with standard cyclically ordered side length collection $(d_0,\dots,d_{n-1}) \in (\mathbb{R}^+)^n$, cyclically re-ordered so that d_0 is largest, if and only if:*

$$\sinh(d_0/2) < \sum_{i=1}^{n-1} \sinh(d_i/2)$$

There is a centered cyclic n-gon in $\mathbb{H}^2$ with standard cyclically ordered side length collection $(d_0,\dots,d_{n-1}) \in (\mathbb{R}^+)^n$, again with d_0 largest, if and only if for $A_d(J)$ as in Lemma 1.7,

$$\sum_{i=1}^{n-1} A_{d_i}(d_0/2) > \pi.$$

Each cyclic n-gon is determined up to isometry of $\mathbb{H}^2$ by the equivalence class of its cyclically ordered side length collection under the action $D_{2n} \times \mathbb{R}^n \to \mathbb{R}^n$ by permutation of entries, and up to orientation-preserving isometry by the action of $\mathbb{Z}_n$.

Proof. Lemma 2.4 implies that $\mathcal{D} = \{d_0,\dots,d_{n-1}\}$ is the side length collection of a cyclic polygon P if and only if the first numerical criterion above is satisfied, and of a centered such polygon P if and only if the second holds. Moreover, if $\{d_0,\dots,d_{n-1}\}$ satisfies the first criterion, then P may be arranged with a starting point $x \in \partial P$ so that proceeding from x in a prescribed direction along ∂P, the edges γ_i are encountered in order.

Given Lemma 2.4, let us fix an n-tuple $(d_0,\dots,d_{n-1})$ with largest element d_0, satisfying $\sinh(d_0/2) < \sum_{i=1}^{n-1}\sinh(d_i/2)$, a polygon $P \subset \mathbb{H}^2$ with edge set $\{\gamma_i\}_{i=0}^{n-1}$ such that $\ell(\gamma_i) =$

d_i for each i, and a point $x \in \gamma_0$ such that the γ_i are encountered in increasing order upon proceeding from x in the direction prescribed by the standard orientation on ∂P. In particular, P has standard cyclically oriented side length collection $(d_0, \dots, d_{n-1})$. We prove the uniqueness criterion below.

Suppose first that P' is a cyclic polygon such that there exists an isometry f of $\mathbb{H}^2$ with $f(P') = P$, and let P' have cyclically ordered side length collection $(d'_0, \dots, d'_{n-1})$. After possibly re-indexing by $\tau\colon\ i \mapsto n - i$, we may assume the cyclic ordering is standard; thus when the edges γ'_i with $\ell(\gamma'_i) = d'_i$ inherit the standard orientation from ∂P, $\gamma'_i \cap \gamma'_{i+1}$ is the terminal endpoint of γ'_i for each i. Since f determines a bijection from the edge set of P' to the edge set of P there exists $\rho \in S_n$ defined by $f(\gamma'_i) = \gamma_{\rho(i)}$. Since f preserves edge lengths, it follows that $d'_i = d_{\rho(i)}$.

Since f takes edges of P' that share a vertex to edges of P with the same property, the definition of cyclic ordering implies that for each i, $\rho(i+1) = \rho(i) \pm 1$. Moreover, it is easy to see that because f takes distinct edges of P' to distinct edges of P, the choice of "+" or "−" above is uniform over all i. If $\rho(i+1) = \rho(i) + 1$ then $\rho = \sigma^k \in \mathbb{Z}_n$, where $k = \rho(0)$, and otherwise $\tau\rho \in \mathbb{Z}_n$.

If f is orientation-preserving on $\mathbb{H}^2$, then in particular it takes the standard orientation on $\partial P'$ to the standard orientation on ∂P. Since the cyclic orderings on the edge sets of P and P' are also standard, in this case $\rho(i+1) = \rho(i) + 1$ and $\rho \in \mathbb{Z}_n$. This proves one direction of the uniqueness criterion.

Now suppose P' has standard cyclically ordered side length collection $(d'_0, \dots, d'_{n-1})$, and for each i $d'_i = d_{\rho(i)}$ for some $\rho \in D_{2n}$. After re-ordering $(d_0, \dots, d_{n-1})$ by σ^k, and $(d'_0, \dots, d'_{n-1})$ by $\sigma^{k'}$, for some $k, k' \in \{0, 1, \dots, n-1\}$, and replacing ρ by $\sigma^{-k}\rho\sigma^{k'}$, we may assume that d_0 and d'_0 are maximal entries of their respective tuples and $\rho(0) = 0$. Therefore ρ is the identity or $\tau\colon\ i \mapsto i + 1$, since these are the only elements of D_{2n} that fix 0. In either case,

$$\sum_{i=1}^{n-1} A_{d'_i}(d'_0/2) = \sum_{i=1}^{n-1} A_{d_{\rho(i)}}(d_0/2) = \sum_{i=1}^{n-1} A_{d_i}(d_0/2)$$

Therefore Lemma 2.4 implies that P and P' are simultaneously centered or non-centered.

First suppose P is centered (and hence also P'), and let J_0 be the radius of P. Then by Lemma 1.10:

$$\sum_{i=0}^{n-1} A_{d'_i}(J) = \sum_{i=0}^{n-1} A_{d_{\rho(i)}}(J) = \sum_{i=0}^{n-1} A_{d_i}(J) = 2\pi.$$

Lemma 2.2 therefore implies that P' also has radius J_0.

Let $\{\gamma_0, \dots, \gamma_{n-1}\}$ and $\{\gamma'_0, \dots, \gamma'_{n-1}\}$ be the cyclically ordered collections of sides of P and P', respectively, so that $\ell(\gamma_i) = d_i$ and $\ell(\gamma'_i) = d'_i$ for each i, and let v and v' be respective centers of P and P'. Decompose P and P' by collections of isosceles triangles $\{T_i\}_{i=0}^{n-1}$ and $\{T'_i\}_{i=0}^{n-1}$, respectively, using the construction from Lemma 1.6.

Since T'_0 and T_0 have identical side length collections they are isometric, and since T_0 is isosceles it admits a self-isometry exchanging the equal-length sides. Therefore there exists an isometry $f\colon \mathbb{H}^2 \to \mathbb{H}^2$ taking T_0' to T_0, so that $T_0' \cap T_1' \mapsto T_0 \cap T_1$ if ρ is the identity, or $T_0' \cap T_1' \mapsto T_0 \cap T_{n-1}$ if $\rho = \tau$.

If ρ is the identity then by the definition above $f(T_1')$ is an isosceles triangle with two sides of length J and one of length $d_1' = d_{\rho(1)} = d_1$ that intersects $f(T_0') = T_0$ in precisely $T_0 \cap T_1$; thus $f(T_1') = T_1$. On the other hand if $\rho = \tau$ then by construction $f(T_1')$ has one side of length $d_1' = d_{\rho(1)} = d_{n-1}$ and intersects $f(T_0') = T_0$ in $T_0 \cap T_{n-1}$, and it follows that $f(T_1') = T_{n-1}$. Arguing inductively we find in the first case that $f(T_i') = T_i$ for each $i > 0$, and in the second that $f(T_i') = T_{n-i}$ for each $i > 0$.

In either case, since $P' = \bigcup_{i=0}^{n-1} T_i'$ and $P = \bigcup_{i=0}^{n-1} T_i$ we have $f(P') = P$. If ρ is the identity then f preserves the standard orientation on $\partial P'$, whereas if $\rho = \tau$ then f reverses it (recall that these were chosen so that γ_0' and γ_0 have respective terminal endpoints $\gamma_0' \cap \gamma_1'$ and $\gamma_0 \cap \gamma_1$). It follows that f preserves orientation if and only if $\rho \in \mathbb{Z}_n$.

If P is non-centered and has radius J_1 then by Lemma 1.10:

$$\sum_{i=1}^{n-1} A_{d_i'}(J_1) = \sum_{i=1}^{n-1} A_{d_{\rho(i)}}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1) = A_{d_0}(J_1) = A_{d_0'}(J_1)$$

Thus P' also has radius J_1. We again produce triangle collections $\{T_i\}$ and $\{T_i'\}$ using the construction from Lemma 1.6. In this case, however, we define an isometry f taking T_1' to T_1 such that $f(T_1' \cap T_2') = T_1 \cap T_2$ if ρ is the identity, and define f to take T_1' to T_{n-1} otherwise, with $f(T_1' \cap T_2) = T_{n-1} \cap T_{n-2}$ in this case.

In the first case it follows by induction that $f(T_i') = T_i$, and in the second that $f(T_i') = T_{n-i}$, for each $i > 1$. Therefore by Lemma 1.6, f takes $P' \cup T_0' = \bigcup_{i=1}^{n-1} T_i'$ to $P \cup T_0 = \bigcup_{i=1}^{n-1} T_i$. Furthermore, one finds that f takes the vertex set of T_0' to that of T_0, so $f(T_0') = T_0$ and it follows that $f(P') = P$. The orientation preserving/reversing dichotomy matches that of the centered case, and the proposition is proved. □

Given a point in $p \in \mathbb{H}^2$ and a one-dimensional subspace $V \subset T_p\mathbb{H}^2$, there are two horocycles through p with tangent space V, one in each half-space bounded by the geodesic γ through p with $V = T_p\gamma$. For such a half-space $\mathcal{H}$, the horocycle in $\mathcal{H}$ is the natural "limit" object for a family of circles through p, each contained in $\mathcal{H}$, whose radii increase without bound.

Making the description above precise would take us too far afield; we only intend it to motivate the relevance of horocyclic polygons (defined below) to the study of cyclic polygons. In proving existence, we will work with an explicitly described horocycle in a model of $\mathbb{H}^2$.

Definition 2.8. A compact, convex polygon in $\mathbb{H}^2$ is *horocyclic* if a single horocycle contains all of its vertices.

The result for horocyclic n-gons that corresponds to Proposition 2.7 differs from it by two important features: each horocyclic n-gon has a unique longest side, and its isometry class is determined by the lengths of the other $n-1$ sides.

Proposition 2.9. *For $n \geq 3$ and each $(d_1, \ldots, d_{n-1}) \in (\mathbb{R}^+)^{n-1}$, there is a horocyclic n-gon P in $\mathbb{H}^2$ with cyclically ordered side length collection $(H_0, d_1 \ldots, d_{n-1})$, where H_0 satisfies:*

$$\sinh(H_0/2) = \sum_{i=1}^{n-1} \sinh(d_i/2)$$

A horocyclic n-gon P' with cyclically ordered side length collection $(H_0', d_1', \ldots, d_{n-1}')$, cyclically re-ordered so that H_0' is largest, is isometric to P by an orientation-preserving map if and only if $d_i' = d_i$ for all i, and isometric to P if and only if $d_i' = d_i$ or $d_i' = d_{n-i}$ for all i.

Proof. We will work in the upper half-plane model for $\mathbb{H}^2$: $\{(x,t) \,|\, t > 0\} \subset \mathbb{R}^2$, and fix the "standard" horocycle $C_\infty = \{(x,1)\}$. The hyperbolic Riemannian metric on $\mathbb{H}^2$ restricts on C_∞ to the Euclidean metric that it inherits from $\mathbb{R}$ under the diffeomorphism $x \mapsto (x,1)$. It is a key fact that if x and y in C have Euclidean distance ℓ, then their hyperbolic distance d satisfies $\sinh(d/2) = \ell/2$.

For fixed $(d_1, \ldots, d_{n-1}) \in (\mathbb{R}^+)^{n-1}$, let $\ell_i = 2\sinh(d_i/2)$ for each i. Define $x_0 = 0$ and $x_i = \sum_{j=1}^{i} \ell_j$ for $i > 0$, and note that the key fact implies that $(x_i, 1)$ and $(x_{i-1}, 1)$ have hyperbolic distance d_i for each $i > 0$. Let H_0 be the hyperbolic distance between $(x_0, 1)$ and $(x_{n-1}, 1)$. Since the Euclidean distance in C_∞ between $(x_0, 1)$ and $(x_{n-1}, 1)$ is $\sum_{i=1}^{n-1} \ell_i$, the key fact gives:

$$2\sinh(H_0/2) = \sum_{i=1}^{n-1} \ell_i = \sum_{i=1}^{n-1} 2\sinh(d_i/2)$$

Hence $H_0 = d((x_0, 1), (x_{n-1}, 1))$ satisfies the equation written in the statement of the lemma.

For each i, the geodesic δ_i containing $(x_{i-1}, 1)$ and $(x_i, 1)$ is the intersection with $\mathbb{H}^2$ of a circle C_i in $\mathbb{R}^2$ centered at $((x_{i-1} + x_i)/2, 0)$, and with radius $r_i = \sqrt{(x_{i-1} - x_i)^2/4 + 1}$. For each $i \in \{0, \ldots, n-1\}$, let $\gamma_i = \{(x,t) \in \delta_i \,|\, t \geq 1\}$ be the geodesic arc joining x_{i-1} to x_i. Further let D_i be the closed disk in $\mathbb{R}^2$ with boundary C_i, let $\mathcal{H}_i = D_i \cap \mathbb{H}^2$, and let $\overline{\mathcal{H}}_i = \overline{\mathbb{H}^2 - \mathcal{H}_i}$. Then $P = \mathcal{H}_0 \cap \left(\bigcap_{i=1}^{n-1} \overline{\mathcal{H}}_i \right)$ is a convex polygon with vertex set $\{(x_i, 1)\}_{i=0}^{n-1}$, sides γ_i, and associated cyclically ordered side length collection $(H_0, \ldots, d_{n-1})$. This order is standard.

Given a horocyclic n-gon P', let $\gamma_0', \ldots, \gamma_{n-1}'$ be the collection of sides of P', cyclically ordered so that $H_0' = \ell(\gamma_0)$ is at least as large as $d_i' = \ell(\gamma_i)$ for each $i > 0$. After possibly re-indexing by $\tau \colon i \mapsto n - i$, we may assume the cyclic order is standard.

If an isometry f of $\mathbb{H}^2$ carries P' to P, then since γ_0' and γ_0 are the unique longest sides of P' and P, respectively, it follows that $H_0' = H_0$ and $f(\gamma_0') = \gamma_0$. Since f preserves adjacency of sides, $f(\gamma_1')$ is either γ_1 or γ_{n-1}. Moreover, if f preserves orientation on $\mathbb{H}^2$ then it takes the standard orientation on $\partial P'$ to the same on ∂P, so since the side length collections are standard it follows in this case that $f(\gamma_1') = \gamma_1$.

It further follows, in the orientation-preserving case, that $f(\gamma_2')$ is adjacent to γ_1 but not equal to γ_0 (since f is one-to-one on the collection of sides of P'), so $f(\gamma_2') = \gamma_2$. Arguing by induction shows in this case that $f(\gamma_i') = \gamma_i$, and hence that $d_i' = d_i$, for all i. On the other hand, if f reverses orientation then $f(\gamma'_1) = \gamma_{n-1}$. An analogous argument shows in this case that $f(\gamma'_i) = \gamma_{n-i}$ and $d'_i = d_{n-i}$ for all $i \in \{1, \ldots, n-1\}$.

Now suppose the standard cyclically ordered side length collection of P' satisfies $H'_0 = H_0$, and either $d'_i = d_i$ for each i, or $d'_i = d_{n-i}$ for each i. The orientation-preserving isometry group of $\mathbb{H}^2$ acts transitively on its collection of horocycles ([cite]). Therefore we will apply such a map and assume that the vertices of P lie on C_∞. Below we will pin down the structure of P' and then construct an isometry taking P' to P.

As was the case with P, each side γ'_i of P' is contained in a geodesic $\delta'_i = C'_i \cap \mathbb{H}^2$, where C'_i is a circle centered in $\mathbb{R} \times \{0\}$. For each i let D'_i be the closed disk in $\mathbb{R}^2$ bounded by C'_i, let $\mathcal{H}'_i = D'_i \cap \mathbb{H}^2$, and let $\overline{\mathcal{H}}'_i = \overline{\mathbb{H}^2 - D'_i}$.

Claim 2.9.1. $P' = \mathcal{H}'_0 \cap \left(\bigcap_{i=1}^{n-1} \overline{\mathcal{H}}'_i \right)$.

Proof of claim. For $i \in \{0, 1, \ldots, n-1\}$, let $x_i \in \mathbb{R}$ be defined by the equation $(x_i, 1) = \gamma_i \cap \gamma_{i+1} = (x_{i+1}, 1)$, taking $i+1$ modulo n. Then $\mathcal{H}'_i \cap C_\infty = I_i \times \{1\}$ is compact, where I_i is the closed interval bounded by x_{i-1} and x_i (either $[x_{i-1}, x_i]$ or $[x_i, x_{i-1}]$, depending on which is larger). Conversely, $\overline{\mathcal{H}}'_i \cap C_\infty = \overline{\mathbb{R} - I_i} \times \{1\}$ is non-compact for each i.

If $P \subset \mathcal{H}'_i$ for some i, then for each $j \neq i$ the endpoints $(x'_j, 1)$ and $(x'_{j-1}, 1)$ of γ'_j are contained in $\mathcal{H}'_i \cap C_\infty = I_i \times \{1\}$, so $I_j \subset I_i$. This containment is proper, since γ_i and γ_j are distinct. It follows that x_i and x_{i-1} are in $\overline{\mathbb{R} - I_j}$ but not I_j, and hence that $P \subset \overline{\mathcal{H}}'_j$. Therefore $P \subset \mathcal{H}'_i$for at most one $i \in \{0, \ldots, n-1\}$.

On the other hand if $P = \bigcap_{i=0}^{n-1} \overline{\mathcal{H}}'_i$, then it would follow that there exists $T > 0$ such that $(x, t) \in P$ as long as $t \geq T$. This would contradict compactness of P, so $P \subset \mathcal{H}'_{i_0}$ for a unique $i_0 \in \{0, \ldots, n-1\}$. The key fact above implies that $2 \sinh \ell(\gamma'_i) = \ell(I_i) = |x_i - x_{i-1}|$ for each i, so since the hyperbolic sine is increasing γ_{i_0} has maximal length among the γ_i. Therefore since we have ordered so that γ'_0 is longest, $i_0 = 0$ and the claim follows. ☐

The standard orientation on $\mathbb{R}^2$ orients C'_0 counterclockwise as the boundary of D'_0. Since $P \subset \mathcal{H}'_0$ the standard orientation on $\gamma'_0 = \partial P \cap C'_0$ matches this orientation, so since the cyclic order on $\{\gamma'_i\}$ was chosen to be standard $x'_0 < x'_{n-1}$, where $(x_i, 1) = \gamma_i \cap \gamma'_{i+1}$. Thus $x'_1 \in (x'_0, x'_{n-1})$, and since $\mathcal{H}'_1 \cap C_\infty$ is non-compact we note that $(x'_0, x'_1) \times \{1\} \cap P' = \emptyset$. Therefore $x'_2 \in (x'_1, x'_{n-1})$, and arguing by induction one establishes:

$$x'_0 < x'_1 < x_{2'} \ldots < x'_{n-1}$$

Now suppose that $d'_i = d_i$ for all i. Then by the key fact it is also true that $|x'_i - x'_{i-1}| = |x_i - x_{i-1}|$ for all i. The map, $(x, t) \mapsto (x - x'_0, t)$ is an orientation-preserving isometry of the upper half-plane model that preserves C_∞ and takes $(x'_0, 1)$ to $(0, 1) = (x_0, 1)$. Upon applying this map, since $x'_1 - x'_0 = x_1 - x_0$ it follows that $x'_1 = x_1$, and similarly by induction that $x'_i = x_i$ for all i.

If $d'_i = d_{n-i}$ for each i, let us first apply the orientation-reversing isometry $(x,t) \mapsto (-x,t)$ of the upper half-plane model that preserves C_∞. Afterwards we have

$$x'_{n-1} < x'_{n-2} < \ldots, x'_1 < x'_0,$$

but the difference $|x'_i - x'_{i-1}|$ is preserved for each i. The orientation-preserving isometry $(x,t) \mapsto (x - x'_{n-1}, t)$ preserves C_∞ and takes $(x'_{n-1}, 1)$ to $(0,1) = (x_0, 1)$. Since $d'_i = d_{n-i}$ for each i, the key fact implies that $|x'_i - x'_{i-1}| = |x_{n-i} - x_{n-i-1}|$ for each i. In particular, taking $i = n-1$ gives $|x'_{n-1} - x'_{n-2}| = |x_1 - x_0|$, so after applying the two maps above we have $x'_{n-2} = x_1$. Arguing inductively gives $x'_i = x_{n-i-1}$ for each i, taking $n-i-1$ modulo n.

In the $d'_i = d_i$ case we have produced an orientation-preserving isometry, and in the $d'_i = d_{n-i}$ case an orientation-reversing one, taking the vertex set of P' to the vertex set of P. Given our descriptions of P and P', it is clear that such an isometry takes P' to P, and the result is proved. □

3. Parametrizing cyclic n-gons

Given Proposition 2.7, it is natural to parametrize the set of oriented cyclic n-gons by their side lengths. We carry this out below, also distinguishing the centered polygons. In the remainder of the section we will describe the boundary of the set of centered polygons in the set of cyclic polygons, and show that the set of horocyclic polygons, properly parametrized, bounds the set of cyclic polygons.

Definition 3.1. For $n \geq 3$, let $\sigma\colon \mathbb{R}^n \to \mathbb{R}^n$ be given by $\sigma(d_0, \ldots, d_{n-1}) = (d_1, \ldots, d_{n-1}, d_0)$, and refer by $\mathbb{R}^n/\mathbb{Z}_n$ to the quotient by the action of $\mathbb{Z}_n \doteq \langle \sigma \rangle$ and by $[d_0, \ldots, d_{n-1}]$ to the equivalence class in $\mathbb{R}^n/\mathbb{Z}_n$ of $(d_0, \ldots, d_{n-1})$. Define:

$$\widetilde{\mathcal{AC}}_n = \left\{ (d_0, \ldots, d_{n-1}) \in (\mathbb{R}^+)^n \mid \sinh(d_i/2) < \sum_{j \neq i} \sinh(d_j/2) \text{ for each } i \in \{0, \ldots, n-1\} \right\}$$

$$\widetilde{\mathcal{C}}_n = \left\{ (d_0, \ldots, d_{n-1}) \in (\mathbb{R}^+)^n \mid \sum_{i=0}^{n-1} A_{d_i}(D/2) > 2\pi, \text{ where } D = \max\{d_i\}_{i=0}^{n-1} \right\}$$

Let $\mathcal{AC}_n \doteq \widetilde{\mathcal{AC}}_n/\mathbb{Z}_n \subset \mathbb{R}^n/\mathbb{Z}_n$ and $\mathcal{C}_n \doteq \widetilde{\mathcal{C}}_n/\mathbb{Z}_n \subset \mathcal{AC}_n$. We say that a cyclic n-gon $P \subset \mathbb{H}^2$ is *represented* by $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n$ or $[d_0, \ldots, d_{n-1}] \in \mathcal{AC}_n$ if there is a standard cyclic order $\gamma_0, \ldots, \gamma_{n-1}$ on the sides of P, in the sense of Definition 2.6, with $\ell(\gamma_i) = d_i$ for each i.

We regard $\widetilde{\mathcal{AC}}_n$ as the set of *marked*, oriented cyclic n-gons in $\mathbb{H}^2$, marked by a choice of which side to measure first. Lemma 2.3 implies in particular that $\widetilde{\mathcal{C}}_n \subset \widetilde{\mathcal{AC}}_n$. By definition, $\widetilde{\mathcal{AC}}_n$ and $\widetilde{\mathcal{C}}_n$ are preserved by the action of the symmetric group S_n on $\mathbb{R}^n$ by permutation of entries, but Proposition 2.7 implies cyclic n-gons P and P' are isometric if and only if any respective representatives $(d_0, \ldots, d_{n-1})$ and $(d'_0, \ldots, d'_{n-1})$ in $\widetilde{\mathcal{AC}}_n$ are D_{2n}-equivalent, where $D_{2n} = \langle \sigma, \tau \rangle$ as described above Definition 2.6.

Proposition 2.7 further implies that P and P' as above are orientation-preserving isometric if and only if $(d_0, \ldots, d_{n-1})$ is $\langle \sigma \rangle$-equivalent to $(d'_0, \ldots, d'_{n-1})$. Thus $\mathcal{AC}_n$ and $\mathcal{C}_n$ parametrize the set of unmarked, oriented, cyclic and centered n-gons in $\mathbb{H}^2$, respectively. Since $\mathbb{Z}_n$ is normal in D_{2n}, τ determines a map of $\mathbb{R}^n/\mathbb{Z}_n$ preserving $\mathcal{AC}_n$ and $\mathcal{C}_n$, so $\mathcal{AC}_n/\tau$ and $\mathcal{C}_n/\tau$ parametrize unmarked, unoriented, cyclic and centered n-gons.

We give $\widetilde{\mathcal{AC}}_n$ the subspace topology from $\mathbb{R}^n$, and note that $\widetilde{\mathcal{C}}_n$ is thus an open subset. The map $\mathbb{R}^n \to \mathbb{R}^n/\mathbb{Z}_n$ is an orbifold quotient, since the action of $\mathbb{Z}_n$ has fixed points, so the fundamental domain U_n defined below will prove convenient in proofs.

Lemma 3.2. *For $n \geq 3$, let $U_n = \{(d_0, \ldots, d_{n-1}) \,|\, d_0 > d_i > 0 \text{ for all } i > 0\} \subset (\mathbb{R}^+)^n$. If $\overline{U}_n$ is the closure of U_n in $(\mathbb{R}^+)^n$, then $\overline{U}_n - U_n \subset \widetilde{\mathcal{C}}_n$, and $U_n \cap \widetilde{\mathcal{AC}}_n$ is a fundamental domain for the $\mathbb{Z}_n$-action on $\widetilde{\mathcal{AC}}_n$.*

Proof. The closure of U_n in $(\mathbb{R}^+)^n$ consists of points $(d_0, \ldots, d_{n-1})$ with $d_0 \geq d_i > 0$ for all $i > 0$, and its frontier $\overline{U}_n - U_n$ consists of such points with $d_0 = d_{i_0}$ for some $i_0 > 0$. For such $(d_0, \ldots, d_{n-1})$:

$$\sum_{i=1}^{n-1} A_{d_i}(d_0/2) = \pi + \sum_{i_0 \neq i,0} A_{d_i}(d_0/2) > \pi$$

This implies that $(d_0, \ldots, d_{n-1}) \in \mathcal{C}_n$, since $A_{d_0}(d_0/2) = \pi$, and it follows that $\overline{U}_n - U_n \subset \widetilde{\mathcal{C}}_n$. Since $\overline{U}_n = \bigcap_{i>0}\{d_0 \geq d_i\}$ is the intersection of convex half-planes in $(\mathbb{R}^+)^n$, it is in particular connected, and its frontier in $(\mathbb{R}^+)^n$, $\overline{U}_n - U_n = \bigcup_{i_0>0}\{d_0 = d_{i_0} \geq d_i \ \forall\, i\}$ is clearly connected as well. For $(d_0, \ldots, d_{n-1}) \in \overline{U}_n$, if $m = \max\{d_i\}_{i>0}$, then taking

$$(t, (d_0, \ldots, d_{n-1})) \mapsto (d_0 - t \cdot (d_0 - m), d_1, \ldots, d_{n-1}) \qquad (t \in [0,1])$$

determines a deformation retract from $\overline{U}_n$ to its frontier. It is easy to see that this restricts on $\widetilde{\mathcal{AC}}_n$ to a deformation retract to $\overline{U}_n - U_n \subset \widetilde{\mathcal{C}}_n$. Since $\overline{U}_n - U_n$ is connected, so therefore is $\overline{U}_n \cap \widetilde{\mathcal{AC}}_n$.

Since any point of U_n has its first entry as the unique largest, any non-trivial cyclic permutation of entries removes this point from U_n. On the other hand, if $(d_0, \ldots, d_{n-1}) \in (\mathbb{R}^+)^n$ has largest entry d_{i_0} then $\sigma^{-i_0}(d_0, \ldots, d_{n-1}) \in \overline{U}_n$, where $\sigma\colon\, i \mapsto i-1$ generates $\mathbb{Z}_n$. Hence $\overline{U}_n$ is a fundamental domain for the $\mathbb{Z}_n$-action on $\mathbb{R}^n$, and $\overline{U}_n - \widetilde{\mathcal{AC}}_n$ is a fundamental domain for the restriction to $\widetilde{\mathcal{AC}}_n$. ☐

Lemma 3.3. *For $n \geq 3$, the closure of $\widetilde{\mathcal{C}}_n$ in $\widetilde{\mathcal{AC}}_n$ (and in $(\mathbb{R}^+)^n$) is:*

$$\left\{ (d_0, \ldots, d_{n-1}) \in (\mathbb{R}^+)^n \,\middle|\, \sum_{i=0}^{n-1} A_{d_i}(D/2) \geq 2\pi, \ \textit{where } D = \max\{d_i\}_{i=0}^{n-1} \right\}$$

This set projects to the closure $\overline{\mathcal{C}}_n$ of $\mathcal{C}_n$ in $\mathcal{AC}_n$.

Proof. Lemma 2.1 implies that for each $i > 0$, $(d_0, \ldots, d_{n-1}) \mapsto A_{d_i}(d_0/2)$ determines a continuous function on $\overline{U}_n$. Thus since $\overline{U}_n$ is already closed, the definition of $\widetilde{\mathcal{C}}_n$ implies that the closure of $\widetilde{\mathcal{C}}_n \cap \overline{U}_n$ in $(\mathbb{R}^+)^n$ is contained in:

$$\left\{ (d_0, \ldots, d_{n-1}) \,|\, d_0 \geq d_i > 0 \ \forall\, i > 0 \text{ and } \sum_{i=0}^{n-1} A_{d_i}(d_0/2) \geq 2\pi \right\} \tag{3.3.1}$$

We will show first that the set above is equal to the closure of $\widetilde{\mathcal{C}}_n \cap \overline{U}_n$ in $(\mathbb{R}^+)^n$. For this we must show that $(d_0, \dots, d_{n-1}) \in \overline{U}_{n-1}$ with $\sum_{i=0}^{n-1} A_{d_i}(d_0/2) = 2\pi$ is approached by a sequence of elements of $\widetilde{\mathcal{C}}_n$. Fix such a point and let $\epsilon = \min\{d_0 - d_i \,|\, 1 \leq i \leq n-1\}$. Since $\overline{U}_n - U_n \subset \widetilde{\mathcal{C}}_n$ (by Lemma 3.2), $\epsilon > 0$, so for $k \in \mathbb{N}$ we have:

$$\sum_{i=1}^{n-1} A_{d_i}((d_0 - \epsilon/k)/2) > \sum_{i=1}^{n-1} A_{d_i}(d_0/2) = \pi.$$

Thus $(d_0 - \epsilon/k, d_1, \dots, d_{n-1}) \in \widetilde{\mathcal{C}}_n$. Since this sequence approaches $(d_0, d_1, \dots, d_{n-1})$ in $\mathbb{R}^n$ as $k \to \infty$, $(d_0, \dots, d_{n-1})$ is in the closure of $\widetilde{\mathcal{C}}_n$.

Let us also note for $(d_0, \dots, d_{n-1}) \in \overline{U}_n$ that since $A_{d_0}(d_0/2) = \pi$, the condition that $\sum_{i=0}^{n-1} A_{d_i}(d_0/2) \geq 2\pi$ is equivalent to $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) \geq \pi$. Lemma 2.3 thus implies that if this holds then $\sinh(d_0/2) < \sum_{i=1}^{n-1} \sinh(d_i/2)$. It follows that the closure of $\widetilde{\mathcal{C}}_n \cap \overline{U}_n$ in $(\mathbb{R}^+)^n$, from (3.3.1), is contained in $\widetilde{\mathcal{AC}}_n$. For $i \in \{0, \dots, n-1\}$, $\sigma^{-i} \in \mathbb{Z}_n$ translates this to:

$$\left\{ (d_0, \dots, d_{n-1}) \,|\, d_i \geq d_j > 0 \ \forall\ j \neq i \text{ and } \sum_{j=0}^{n-1} A_{d_j}(d_i/2) \geq 2\pi \right\}$$

This set is the closure of $\widetilde{\mathcal{C}}_n \cap \sigma^{-i}(\overline{U}_n)$, and since $\overline{U}_n$ is a fundamental domain for the $\mathbb{Z}_n$-action on $\widetilde{\mathcal{AC}}_n$ it follows that the closure of $\widetilde{\mathcal{C}}_n$ is as claimed. This projects to $\overline{\mathcal{C}}_n$ in $\mathbb{R}^n/\mathbb{Z}_n$. ☐

Lemma 3.4. *For $n \geq 3$, there is a smooth function $b_0 \colon (\mathbb{R}^+)^{n-1} \to \mathbb{R}^+$ such that for each $(d_1, \dots, d_{n-1}) \in (\mathbb{R}^+)^{n-1}$, $J = \frac{1}{2} b_0(d_1, \dots, d_{n-1})$ uniquely solves $\sum_{i=1}^{n-1} A_{d_i}(J) = \pi$. This further satisfies $b_0(d_1, \dots, d_{n-1}) > \max\{d_i\}_{i=1}^{n-1}$, and $b_0(d_1, \dots, d_{n-1}) \leq b_0(d_1', \dots, d_{n-1}')$ whenever $d_i \leq d_i'$ for all i. Define:*

$$\mathrm{Graph}(b_0) = \left\{ (b_0(d_1, \dots, d_{n-1}), d_1, \dots, d_{n-1}) \,|\, (d_1, \dots, d_{n-1}) \in (\mathbb{R}^+)^{n-1} \right\} \subset U_n \cap \widetilde{\mathcal{AC}}_n$$

Then $\widetilde{\mathcal{BC}}_n \doteq \mathbb{Z}_n \cdot \mathrm{Graph}(b_0)$ is the frontier of $\widetilde{\mathcal{C}}_n$ in $\widetilde{\mathcal{AC}}_n$, so $\widetilde{\mathcal{C}}_n \sqcup \widetilde{\mathcal{BC}}_n$ is the closure of $\widetilde{\mathcal{C}}_n$ in $\widetilde{\mathcal{AC}}_n$. Furthermore, $\widetilde{\mathcal{BC}}_n$ projects to the frontier $\mathcal{BC}_n$ of $\mathcal{C}_n$ in $\mathcal{AC}_n$.

Proof. For fixed $(d_1, \dots, d_{n-1}) \in (\mathbb{R}^+)^{n-1}$ with maximal entry d_{i_0}, by Lemma 1.8 the sum $\sum_{i=1}^{n-1} A_{d_i}(J)$ is well-defined and decreasing on $[d_{i_0}/2, \infty)$, taking a value greater than π at $J = d_{i_0}/2$ and with a horizontal asymptote of 0. There is thus a unique B_0, greater than d_i for all i, such that $F(B_0/2, \dots, d_{n-1}) = 0$, and we define $b_0(d_1, \dots, d_{n-1}) = B_0$.

By construction, $\sum_{i=1}^{n-1} A_{d_i}(J) = \pi$ is uniquely satisfied at $J = \frac{1}{2} b_0(d_1, \dots, d_{n-1})$ for each $(d_1, \dots, d_{n-1}) \in (\mathbb{R}^+)^{n-1}$, and also $b_0(d_1, \dots, d_{n-1}) > \max\{d_i\}_{i=1}^{n-1}$. Since $A_d(J)$ increases in d (by Lemma 1.8 again), for $(d_1, \dots, d_{n-1})$ and $(d_1', \dots, d_{n-1}')$ with $d_i \leq d_i'$ for each i we have $\sum_{i=1}^{n-1} A_{d_i'}(\frac{1}{2} b_0(\frac{1}{2} d_1, \dots, d_{n-1}), d_1, \dots, d_{n-1}) \geq \pi$, so $b_0(d_1', \dots, d_{n-1}') \geq b_0(d_1, \dots, d_{n-1})$.

We will use the implicit function theorem to show b_0 is smooth. For U_n as in Lemma 3.2, define $F \colon \overline{U}_n \to \mathbb{R}$ by $F(d_0, \dots, d_{n-1}) = \pi - \sum_{i=1}^{n-1} A_{d_i}(d_0/2)$. Lemma 2.1 implies that F is continuous on $\overline{U}_n$ and smooth on U_n. We recorded the derivative of $A_d(J)$ with respect to

J in (2.1.1); this gives:

$$\frac{\partial F}{\partial d_0} = -\coth(d_0/2)\sum_{i=1}^{n-1}\frac{\sinh(d_i/2)}{\sqrt{\sinh^2(d_0/2)-\sinh^2(d_i/2)}}$$

By construction, $F(d_0,\ldots,d_{n-1}) = 0$ if and only if $d_0 = b_0(d_1,\ldots,d_{n-1})$. At such a point all summands are positive in the formula above; hence $\partial F/\partial d_0 < 0$ here, so the implicit function theorem implies that b_0 is smooth..

Since $A_d(d/2) = \pi$, each $(d_0, d_1, \ldots, d_{n-1}) \in \mathrm{Graph}(b_0)$ satisfies $\sum_{i=0}^{n-1} A_{d_i}(d_0/2) = 2\pi$, so by Lemma 3.3 $\mathrm{Graph}(b_0)$ is contained in the closure of $\widetilde{\mathcal{C}}_n$ in $\widetilde{\mathcal{AC}}_n$, and the same holds true for each of its $\mathbb{Z}_n$-translates. On the other hand, it follows directly from Definition 3.1 that $\widetilde{\mathcal{BC}}_n$, as we have defined it above, does not intersect $\widetilde{\mathcal{C}}_n$; thus it is the frontier of $\widetilde{\mathcal{C}}_n$ in $\widetilde{\mathcal{AC}}_n$. Since $\widetilde{\mathcal{BC}}_n$ is $\mathbb{Z}_n$-invariant by construction, it projects to the frontier of $\mathcal{C}_n$ in $\mathcal{AC}_n$. □

Corollary 3.5. *For $n \geq 3$ and U_n as in Lemma 3.2 , $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$ has n components, each of the form $\sigma^i(U_n) \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$ for $i \in \{0, \ldots, n-1\}$.*

Proof. Let us recall from Lemma 3.2 that $U_n \cap \widetilde{\mathcal{AC}}_n$ is a fundamental domain for the $\mathbb{Z}_n$-action, and that the frontier $\overline{U}_n - U_n$ of U_n in $\mathbb{R}^n$ is entirely contained in the open subset $\widetilde{\mathcal{C}}_n$ of $\widetilde{\mathcal{AC}}_n$. Thus for each $i \in \{0, \ldots, n-1\}$, the the open subset $\sigma^i(U_n) \cap (\widetilde{\mathcal{AC}}_n - \mathcal{C}_n)$ contains at least one component of $\widetilde{\mathcal{AC}}_n - \mathcal{C}_n$.

By Lemma 3.4 and the definitions, $\mathrm{Graph}(b_0) \subset U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$, and thus it follows from the definition of $\widetilde{\mathcal{BC}}_n$ that $\mathrm{Graph}(b_0) = \widetilde{\mathcal{BC}}_n \cap U_n$. Since $\mathrm{Graph}(b_0)$ is the continuous image of the connected set $(\mathbb{R}^+)^{n-1}$ it is also connected.

If $(d_0, \ldots, d_{n-1}) \in U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$, then by the definitions, $\sum_{i=0}^{n-1} A_{d_i}(d_0/2) \leq \pi$. Since the functions A_{d_i} are decreasing (cf. 1.8), it therefore follows from the definition of b_0 in Lemma 3.4 that $d_0 \geq b_0(d_1, \ldots, d_{n-1})$. We construct a deformation retract $U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n) \to U_n \cap \widetilde{\mathcal{BC}}_n$ by:

$$(t, (d_0, \ldots, d_{n-1})) \mapsto (d_0 - t \cdot (d_0 - b_0(d_1, \ldots, d_{n-1})), d_1, \ldots, d_{n-1}) \qquad (t \in [0,1])$$

It follows that $U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$ is connected, and therefore that $\sigma^i(U_n) \cap (\widetilde{\mathcal{AC}}_n - \mathcal{C}_n)$ is also connected for each $i \in \{0, \ldots, n-1\}$. □

Below we re-interpret earlier work to define a "radius function" on $\widetilde{\mathcal{AC}}_n$. By a *symmetric* function of n-variables, we refer to one that takes the same value on any two n-tuples related by permuting entries. (We noted above that $\widetilde{\mathcal{AC}}_n$ is invariant under such permutations.)

Lemma 3.6. *For $n \geq 3$ and $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n$, let $J(d_0, \ldots, d_{n-1}) = J \geq \max\{d_i/2\}_{i=0}^{n-1}$, where J solves one of the equations below:*

$$\begin{cases} A_{d_{i_0}}(J) = \sum_{i \neq i_0} A_{d_i}(J) & \textit{where } d_{i_0} \geq d_i \ \forall\ i; \ \textit{or} \\ \sum_{i=0}^{n-1} A_{d_i}(J) = 2\pi \end{cases}$$

Then $J(d_0, \ldots, d_{n-1})$ is the radius of the cyclic n-gon represented by $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n$ (see Definition 3.1). It is symmetric on $\widetilde{\mathcal{AC}}_n$ and determines $J\colon \mathcal{AC}_n \to \mathbb{R}^+$.

Proof. By Lemma 2.2, a unique $J \geq D/2$ solves one of the equations above, where $D = \max\{d_i\}_{i=0}^{n-1}$. Thus $J(d_0, \ldots, d_{n-1})$ is a well-defined function on $\widetilde{\mathcal{AC}}_n$. Furthermore, J solves the lower equation above if and only if $\sum_{i=0}^{n-1} A_{d_i}(D/2) \geq 2\pi$, since $A_d(d/2) = \pi$ by Lemma 1.8. For any fixed $J \geq D/2$, the value $\sum_{i=0}^{n-1} A_{d_i}(J)$ is invariant under permutation of the d_i, as is D itself. It follows that $J(d_0, \ldots, d_{n-1})$ is invariant under the S_n-action on $\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$, which by Lemmas 3.4 and 3.3 is the set of $(d_0, \ldots, d_{n-1})$ such that $\sum A_{d_i}(D/2) \geq 2\pi$.

For other $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n$, if $d_i' = d_{\pi(i)}$ for each i, where π is a permutation, then for $d_{i_0} = D$ as above $d'_{\pi^{-1}(i_0)} = D$, and for $J = J(d_0, \ldots, d_{n-1})$ we have:

$$A_{d'_{\pi^{-1}(i_0)}}(J) = A_{d_{i_0}}(J) = \sum_{i \neq i_0} A_{d_i}(J) = \sum_{i \neq \pi^{-1}(i_0)} A_{d_i'}(J)$$

This is because $A_{d_i'}(J) = A_{d_{\pi(i)}}(J)$ for each i, and because $\pi(i) \neq i_0$ for $i \neq \pi^{-1}(i_0)$. Therefore J is symmetric, and in particular it induces a well-defined function on $\mathcal{AC}_n$.

Since $J(d_0, \ldots, d_{n-1})$ uniquely solves one of the equations above, Lemma 1.10 implies that it is the radius of the cyclic n-gon determined by $(d_0, \ldots, d_{n-1})$ (see also Corollary 1.11). $\square$

Since the radius reflects only the intrinsic geometry of a cyclic polygon, one expects that J should not depend on the choice of marking and should determine a well-defined function on $\mathcal{AC}_n$. The functions of the lemma below, on the other hand, reflect geometric quantities associated to a particular side or vertex of a cyclic polygon. Hence we expect that marking-invariance should not hold; it is replaced by a "marking-equivariance" property.

Lemma 3.7. *For $n \geq 3$, $i \in \{0, \ldots, n-1\}$, $J\colon \widetilde{\mathcal{AC}}_n \to \mathbb{R}^+$ as in Lemma 3.6, and $A_d(J)$ and $B_d(J)$ as in Lemma 1.7, let $\alpha_i\colon \widetilde{\mathcal{AC}}_n \to (0, \pi]$ and $\beta_i\colon \widetilde{\mathcal{AC}}_n \to [0, \pi/2)$ be given by:*

$$\alpha_i(d_0, \ldots, d_{n-1}) = A_{d_i}(J(d_0, \ldots, d_{n-1})) \qquad \beta_i(d_0, \ldots, d_{n-1}) = B_{d_i}(J(d_0, \ldots, d_{n-1}))$$

For each i and j, $\alpha_i = \alpha_{i-j} \circ \sigma^j$ and $\beta_i = \beta_{i-j} \circ \sigma^j$, where $i - j$ is taken modulo n. Also:

- *For each i, the relation $\alpha_i(d_0, \ldots, d_{n-1}) = \sum_{j \neq i} \alpha_j(d_0, \ldots, d_{n-1})$ holds if and only if $(d_0, \ldots, d_{n-1}) \in \sigma^{-i}(\overline{U}_n) \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$; and*
- *$2\pi = \sum_{i=0}^{n-1} \alpha_i(d_0, \ldots, d_{n-1})$ if and only if $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$.*

If P is represented by $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n$, with center v, then T_i from Lemma 1.6 has angle $\alpha_i(d_0, \ldots, d_{n-1})$ at v. For each i, β_i satisfies $\sin\beta_i \sinh d_i \equiv \sin\alpha_i \sinh J$ on $\widetilde{\mathcal{AC}}_n$.

Proof. That $\alpha_i(d_0, \ldots, d_{n-1})$ records the angle of T_i follows from its definition, Lemma 3.6, and Lemma 1.10. The relations $\alpha_i = \alpha_{i-j} \circ \sigma^j$ hold because J is $\mathbb{Z}_n$-invariant and we have:

$$\sigma^j(d_0, \ldots, d_{n-1}) = (d_j, \ldots, d_{n-1}, d_0, \ldots, d_{j-1})$$

In particular, the $(i-j)$th coordinate of σ^j is d_i. For the same reason, $\beta_i = \beta_{i-j} \circ \sigma^j$.

The relation $\sin\beta_i \sinh d_i \equiv \sin\alpha_i \sinh J$ can be verified directly from the definitions in Lemma 1.7 with some hyperbolic trigonometry, or by appealing to the geometric interpretations there and the hyperbolic law of sines.

Fix $(d_0,\dots,d_{n-1}) \in \overline{U}_n \cap \widetilde{\mathcal{AC}}_n$; that is, with $d_0 \geq d_i > 0$ for each $i > 0$ and $\sinh(d_0/2) < \sum_{i=1}^{n-1}\sinh(d_i/2)$. Since $\alpha_i(d_0,\dots,d_{n-1}) = A_{d_i}(J(d_0,\dots,d_{n-1}))$, Lemma 3.6 implies that at least one of the following holds:

$$2\pi = \sum_{i=0}^{n-1}\alpha_i(d_0,\dots,d_{n-1}) \qquad\qquad \alpha_0(d_0,\dots,d_{n-1}) = \sum_{i=1}^{n-1}\alpha_i(d_0,\dots,d_{n-1})$$

Lemma 2.2 implies that the identity on the right holds if and only if $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) \leq \pi$; ie, if and only if $(d_0,\dots,d_{n-1}) \in \overline{U}_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$ by definition. The left-hand identity holds if and only if $\sum_{i=1}^{n-1} A_{d_i}(d_0/2) \geq \pi$; ie, if and only if $\sum_{i=0}^{n-1} A_{d_i}(d_0/2) \geq 2\pi$. Lemma 3.3 implies that this holds if and only if $(d_0,\dots,d_{n-1})$ is in the closure of $\widetilde{\mathcal{C}}_n$ in $\widetilde{\mathcal{AC}}_n$. Therefore by Lemma 3.4, $2\pi = \sum_{i=0}^{n-1}\alpha_i(d_0,\dots,d_{n-1})$ if and only if $(d_0,\dots,d_{n-1}) \in \overline{U}_n \cap (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n)$.

For $(d_0,\dots,d_{n-1}) \in \sigma^{-i}(\overline{U}_n) \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$, if $(d_0',\dots,d_{n-1}') = \sigma^i(d_0,\dots,d_{n-1}) \in \overline{U}_n$ then $\alpha_j(d_0,\dots,d_{n-1}) = \alpha_{j-i}(d_0',\dots,d_{n-1}')$ for each j, by Lemma 3.7. In particular:

$$\alpha_i(d_0,\dots,d_{n-1}) = \alpha_0(d_0',\dots,d_{n-1}') \quad \text{and} \quad \sum_{j\neq i}\alpha_j(d_0,\dots,d_{n-1}) = \sum_{j\neq 0}\alpha_j(d_0',\dots,d_{n-1}')$$

Applying the case above, we find that these quantities are equal if and only if $(d_0',\dots,d_{n-1}') \in \overline{U}_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$. Since $\widetilde{\mathcal{C}}_n$ and $\widetilde{\mathcal{AC}}_n$ are σ-invariant, this holds if and only if $(d_0,\dots,d_{n-1}) \in \sigma^{-i}(\overline{U}_n) \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$. Since $\widetilde{\mathcal{BC}}_n$ is also σ-invariant, the same argument shows that $2\pi = \sum_{i=0}^{n-1}\alpha_i(d_0,\dots,d_{n-1})$ if and only if $(d_0,\dots,d_{n-1}) \in \sigma^{-i}(\overline{U}_n) \cap (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n)$. □

Lemma 3.8. *For $n \geq 3$ and $(d_0,\dots,d_{n-1}) \in \widetilde{\mathcal{C}}_n \cup U_n$ (where U_n is as in Lemma 3.2), define $\nu_i(d_0,\dots,d_{n-1}) = (\beta_i+\beta_{i+1})(d_0,\dots,d_{n-1}) \in [0,\pi)$ for $0 < i < n-1$; and let:*

$$\nu_0(d_0,\dots,d_{n-1}) = \begin{cases} (\beta_0+\beta_1)(d_0,\dots,d_{n-1}) & (d_0,\dots,d_{n-1}) \in \widetilde{\mathcal{C}}_n \\ (\beta_1-\beta_0)(d_0,\dots,d_{n-1}) & (d_0,\dots,d_{n-1}) \in U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n) \end{cases}$$

and

$$\nu_{n-1}(d_0,\dots,d_{n-1}) = \begin{cases} (\beta_0+\beta_{n-1})(d_0,\dots,d_{n-1}) & (d_0,\dots,d_{n-1}) \in \widetilde{\mathcal{C}}_n \\ (\beta_{n-1}-\beta_0)(d_0,\dots,d_{n-1}) & (d_0,\dots,d_{n-1}) \in U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n) \end{cases}$$

For each $i \in \{0,\dots,n-1\}$, defining $\nu_i|_{\sigma^{-j}(\overline{U}_n)} = \nu_{i-j}|_{\overline{U}_n} \circ \sigma^j$ determines a well-defined extension to $\widetilde{\mathcal{AC}}_n$ satisfying $\nu_i = \nu_{i-j}\circ\sigma^j$. If P is represented by $(d_0,\dots,d_{n-1}) \in \widetilde{\mathcal{AC}}_n$ (see Definition 3.1), then $\nu_i(d_0,\dots,d_{n-1}) = \angle_{x_i}P$ for each i, where $x_i = \gamma_i \cap \gamma_{i+1}$.

Proof. This is a direct consequence of Lemma 1.10 and the definitions above. □

The lemma below characterizes $\widetilde{\mathcal{BC}}_n$ as a locus of values of the radius function or the angle functions α_i or β_i, and also by a geometric property of the associated polygons.

Lemma 3.9. *For $n \geq 3$, the functions α_i, β_i, and J on $\widetilde{\mathcal{AC}}_n$ satisfy:*

- *$\alpha_i(d_0,\dots,d_{n-1}) = \pi$ if and only if $(d_0,\dots,d_{n-1}) \in \sigma^{-i}(U_n) \cap \widetilde{\mathcal{BC}}_n = \sigma^{-i}(\mathrm{Graph}(b_0))$;*
- *$\beta_i(d_0,\dots,d_{n-1}) = 0$ if and only if $(d_0,\dots,d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$; and*
- *$J(d_0,\dots,d_{n-1}) = d_i/2$ if and only if $(d_0,\dots,d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$.*

Furthermore, $(d_0,\ldots,d_{n-1}) \in \widetilde{\mathcal{AC}}_n$ *is in* $\widetilde{\mathcal{BC}}_n$ *if and only if the cyclic polygon* P *that it determines has its center* v *in* ∂P*; hence* $(d_0,\ldots,d_{n-1}) \in \widetilde{\mathcal{C}}_n \sqcup \widetilde{\mathcal{BC}}_n$ *if and only if* $v \in P$*.*

Proof. For some $(d_0,\ldots,d_{n-1}) \in \widetilde{\mathcal{AC}}_n$, suppose $\alpha_0(d_0,\ldots,d_{n-1}) = \pi$. We note first that for $i \neq 0$, $(d_0,\ldots,d_{n-1}) \notin \sigma^{-i}(\overline{U}_n) \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$: if this were the case for some i, then by Lemma 3.7 the identity below would hold:

$$\alpha_i(d_0,\ldots,d_{n-1}) = \sum_{j\neq i} \alpha_j(d_0,\ldots,d_{n-1})$$

Since $\alpha_j(d_0,\ldots,d_{n-1}) > 0$ for $j \neq i, 0$, this contradicts the fact that $\alpha_i(d_0,\ldots,d_{n-1}) \leq \pi$.

Thus if $\alpha_0(d_0,\ldots,d_{n-1}) = \pi$ then $(d_0,\ldots,d_{n-1}) \in \widetilde{\mathcal{C}}_n$ or $(d_0,\ldots,d_{n-1}) \in \overline{U}_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$. Therefore by Lemma 3.7, the α_i satisfy at least one of the following:

$$2\pi = \sum_{i=0}^{n-1} \alpha_i(d_0,\ldots,d_{n-1}) \qquad\qquad \alpha_0(d_0,\ldots,d_{n-1}) = \sum_{i=1}^{n-1} \alpha_i(d_0,\ldots,d_{n-1})$$

But since $\alpha_0(d_0,\ldots,d_{n-1}) = \pi$, either of the equations above implies the other, so both hold. Lemma 3.7 implies that the locus where both of these equations hold is

$$\overline{U}_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n) \cap (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n) = \overline{U}_n \cap \widetilde{\mathcal{BC}}_n = \mathrm{Graph}(b_0)$$

Thus if $(d_0,\ldots,d_{n-1})$ satisfies $\alpha(d_0,\ldots,d_{n-1}) = \pi$ then $(d_0,\ldots,d_{n-1}) \in \mathrm{Graph}(b_0)$. On the other hand, if $(d_0,\ldots,d_{n-1}) \in \mathrm{Graph}(b_0)$ then since both equations above hold it follows easily that $\alpha_0(d_0,\ldots,d_{n-1}) = \pi$.

The above and Lemma 3.7 imply for each i that $\alpha_i(d_0,\ldots,d_{n-1}) = \alpha_0 \circ \sigma^i(d_0,\ldots,d_{n-1}) = \pi$ if and only if $(d_0,\ldots,d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$. Since $\alpha_i(d_0,\ldots,d_{n-1}) = A_{d_i}(J(d_0,\ldots,d_{n-1}))$ by definition, Lemma 1.8 implies for each i that:

$$J(d_0,\ldots,d_{n-1}) = d_i/2 \quad\Leftrightarrow\quad \alpha_i(d_0,\ldots,d_{n-1}) = \pi \quad\Leftrightarrow\quad (d_0,\ldots,d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$$

Since $d_i > 0$, $J(d_0,\ldots,d_{n-1}) > 0$, and $\alpha_i(d_0,\ldots,d_{n-1}) \in (0,\pi]$, the identity $\sin\beta_i \sinh d_i = \sin\alpha_i \sinh J$ from Lemma 3.7 implies:

$$\beta_i(d_0,\ldots,d_{n-1}) = 0 \quad\Leftrightarrow\quad \alpha_i(d_0,\ldots,d_{n-1}) = \pi \quad\Leftrightarrow\quad (d_0,\ldots,d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$$

Now let P be a cyclic polygon with center v and cyclically ordered side length collection $d_0,\ldots,d_{n-1}$. It is clear that the radius of P is $d_i/2$ if and only if v is the midpoint of a side with length d_i. Lemma 3.6 implies that $J(d_0,\ldots,d_{n-1})$ is the radius of P, so the above gives $(d_0,\ldots,d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$ if and only if v is the center of a side with length d_i. Since $\widetilde{\mathcal{BC}}_n = \bigcup_i \sigma^{-i}(\mathrm{Graph}(b_0))$, it follows that $(d_0,\ldots,d_{n-1}) \in \widetilde{\mathcal{BC}}_n$ if and only if $v \in \partial P$. □

We will finish this section by parametrizing the set of horocyclic n-gons.

Lemma 3.10. *For $n \geq 3$, let $h_0(d_1, \ldots, d_{n-1}) = 2\sinh^{-1}\left(\sum_{i=1}^{n-1} \sinh(d_i/2)\right)$. Then h_0 is smooth on $(\mathbb{R}^+)^{n-1}$, and if $d_i \leq d_i'$ for all i then $h_0(d_1, \ldots, d_{n-1}) \leq h_0(d_1', \ldots, d_{n-1}')$. Define:*

$$\begin{aligned} \mathrm{Graph}(h_0) &= \left\{(h_0(d_1, \ldots, d_{n-1}), d_1, \ldots, d_{n-1}) \,|\, (d_1, \ldots, d_{n-1}) \in (\mathbb{R}^+)^{n-1}\right\} \\ &= \left\{ (H_0, d_1, \ldots, d_{n-1}) \,|\, d_i > 0 \ \forall\ i \text{ and } \sinh(H_0/2) = \sum_{i=1}^{n-1} \sinh(d_i/2) \right\} \subset U_n \end{aligned}$$

$\widetilde{\mathcal{HC}}_n \doteq \mathbb{Z}_n \cdot \mathrm{Graph}(h_0)$ is in one-one correspondence with the set of marked horocyclic n-gons. It is the frontier of $\widetilde{\mathcal{AC}}_n$ in $(\mathbb{R}^+)^n$, and its quotient $\mathcal{HC}_n \in (\mathbb{R}^+)^n/\mathbb{Z}_n$ is the frontier of $\mathcal{AC}_n$.

This lemma is a straightforward consequence of Proposition 2.9 and Definition 3.1. Below we record a useful description of "vertical rays" in $\widetilde{\mathcal{AC}}_n$.

Corollary 3.11. *For $n \geq 3$ and $(d_1, \ldots, d_{n-1}) \in (\mathbb{R}^+)^{n-1}$, $(d_0, d_1, \ldots, d_{n-1}) \in U_n$ is in $\widetilde{\mathcal{C}}_n$ if and only if $d_0 < b_0(d_1, \ldots, d_{n-1})$, and in $\widetilde{\mathcal{AC}}_n$ if and only if $d_0 < h_0(d_1, \ldots, d_{n-1})$, for $b_0 \colon (\mathbb{R}^+)^{n-1} \to \mathbb{R}^+$ as in Lemma 3.4 and h_0 as in Lemma 3.10.*

This is a consequence of Definition 3.1, the definitions of b_0 and h_0, and the fact that $A_d(J)$ is decreasing (by Lemma 1.8), and $\sinh J$ increasing, in J. Also recall that Lemma 2.3 implies that $b_0(d_1, \ldots, d_{n-1}) < h_0(d_1, \ldots, d_{n-1})$.

4. Differentiable dependence on side lengths

The main results of this section show that the radius and vertex angles of cyclic polygons are continuously differentiable functions of their side lengths. It complicates matters that this is not quite true for the central angles of the isosceles triangles in the decomposition from Lemma 1.6: a discontinuity in the derivative of α_i on $\sigma^{-i}(\mathrm{Graph}(b_0))$ (where the corresponding triangle T_i is degenerate) reflects the fact that the interior of T_i "changes sides" as a family of cyclic polygons goes from centered to non-centered.

We also bound partial derivatives of the radius, in Lemma 4.5. At the end of the section we describe the limiting behavior of angles and radius as polygons approach horocyclic.

Proposition 4.1. *For $n \geq 3$, the functions J (from Lemma 3.6) and α_i (from Lemma 3.7), for $i \in \{0, \ldots, n-1\}$, are smooth on $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{BC}}_n$; J is C^1 on all of $\widetilde{\mathcal{AC}}_n$; and α_i is C^0 on $\widetilde{\mathcal{AC}}_n$ and C^1 on $\widetilde{\mathcal{AC}}_n - \sigma^{-i}(\mathrm{Graph}(b_0))$ for each i, where b_0 is as in Lemma 3.4.*

For U_n as in Lemma 3.2 and $i, j \in \{0, \ldots, n-1\}$, we have:

$$\frac{\partial \alpha_i}{\partial d_j} = \begin{cases} -\sum_{k \neq i} \frac{\partial \alpha_k}{\partial d_j} & (d_0, \ldots, d_{n-1}) \in \sigma^{-i}(\overline{U}_n) \cap \widetilde{\mathcal{C}}_n \\ \sum_{k \neq i} \frac{\partial \alpha_k}{\partial d_j} & (d_0, \ldots, d_{n-1}) \in \sigma^{-i}(\overline{U}_n) - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n) \end{cases} \tag{4.1.1}$$

At $(d_0,\dots,d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$, $\frac{\partial\alpha_i}{\partial d_j}$ *has well-defined values computed from within* $\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$ *or* $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$*, and these are opposite. The following identities hold on all of* $\widetilde{\mathcal{AC}}_n$*:*

$$\cot(\alpha_i/2)\frac{\partial\alpha_i}{\partial d_j} + 2\coth J\frac{\partial J}{\partial d_j} = 0 \qquad (i \neq j) \tag{4.1.2}$$

$$\cot(\alpha_j/2)\frac{\partial\alpha_j}{\partial d_j} + 2\coth J\frac{\partial J}{\partial d_j} = \coth(d_j/2) \tag{4.1.3}$$

Proof. Define $f = (\alpha_0,\dots,\alpha_{n-1},J)\colon \widetilde{\mathcal{AC}}_n \to \mathbb{R}^{n+1}$, where J is as in Lemma 3.6 and the α_i are as in Lemma 3.7. We will show smoothness of f, and hence also of J and the α_i, using the implicit function theorem.

Define $F^{(0)}\colon \mathbb{R}^n \times (0,2\pi)^n \times \mathbb{R}^+ \to \mathbb{R}^{n+1}$ by $F^{(0)} = (F^{(0)}_0, F^{(0)}_1, \dots, F^{(0)}_n)$, where:

$$F^{(0)}_i(x_0,\dots,x_{n-1},y_0,\dots,y_{n-1},y_n) = \begin{cases} \sinh^2 y_n - \frac{\cosh x_i - 1}{1-\cos y_i} & 0 \le i < n \\ 2\pi - \sum_{j=0}^{n-1} y_j & i = n \end{cases}$$

$F^{(0)}_i$ is smooth for each i, and for $i<n$, $F^{(0)}_i(d_0,\dots,d_{n-1},\alpha_0,\dots,\alpha_{n-1},J) = 0$ if and only if:

$$\sinh^2 J = \frac{\cosh d_i - 1}{1-\cos\alpha_i} \tag{4.1.4}$$

Comparing with Lemma 1.7, we find that $(d_0,\dots,d_{n-1},\alpha_0,\dots,\alpha_{n-1},J) \in \mathbb{R}^n \times (0,\pi]^n \times \mathbb{R}^+$ is taken by $F^{(0)}_i$ to 0 if and only if $\alpha_i = A_{d_i}(J)$. Thus the definition, in Lemma 3.7, of $\alpha_i\colon \widetilde{\mathcal{AC}}_n \to (0,\pi]$ and the definition above of f imply $F^{(0)}_i(d_0,\dots,d_{n-1},f(d_0,\dots,d_{n-1})) = 0$ for each $i<n$ and $(d_0,\dots,d_{n-1}) \in \widetilde{\mathcal{AC}}_n$.

Lemma 3.7 further implies that $\sum_{i=0}^{n-1}\alpha_i \equiv 2\pi$ on $\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$, so the definition of $F^{(0)}_n$ gives:

$$F^{(0)} \circ (\mathrm{Id}_{\mathbb{R}^n} \times f)|_{\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n} \equiv (0,\dots,0) \tag{4.1.5}$$

However if $(d_0,\dots,d_{n-1}) \notin \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$ it is not necessarily true that $\sum_{i=0}^{n-1}\alpha_i(d_0,\dots,d_{n-1}) = 2\pi$, so we cannot conclude that $(d_0,\dots,d_{n-1})$ is a zero of $F^{(0)} \circ (\mathrm{Id}_{\mathbb{R}^n} \times f)$.

At $(d_0,\dots,d_{n-1},\alpha_0,\dots,\alpha_{n-1},J)$ sent by $F^{(0)}$ to the origin of $\mathbb{R}^{n+1}$, the derivative matrix $dF^{(0)}$ is of the form $(A^{(0)} \mid B^{(0)})$, where $A^{(0)} = \left(\partial F^{(0)}_i/\partial x_j\right)^{0\le i\le n}_{0\le j<n}$ and $B^{(0)} = \left(\partial F^{(0)}_i/\partial y_j\right)^{0\le i\le n}_{0\le j\le n}$ are as below.

$$A^{(0)} = \left(\begin{array}{ccc} \frac{-\sinh d_0}{1-\cos\alpha_0} & & 0 \\ & \ddots & \\ 0 & & \frac{-\sinh d_{n-1}}{1-\cos\alpha_{n-1}} \\ \hline 0 & \dots & 0 \end{array}\right) \quad B^{(0)} = \left(\begin{array}{ccc|c} \frac{\sinh^2 J\sin\alpha_0}{1-\cos\alpha_0} & & 0 & 2\sinh J\cosh J \\ & \ddots & & \vdots \\ 0 & & \frac{\sinh^2 J\sin\alpha_{n-1}}{1-\cos\alpha_{n-1}} & 2\sinh J\cosh J \\ \hline -1 & \dots & -1 & 0 \end{array}\right)$$

To simplify the above we have used the identities (4.1.4) for each i, which imply in particular that $-\frac{\sinh d_i}{1-\cos\alpha_i} = \sinh^2 J \cdot \frac{-\sinh d_i}{\cosh d_i - 1}$.

We claim that $B^{(0)}$ is invertible at such $(d_0,\dots,d_{n-1},\alpha_0,\dots,\alpha_{n-1},J)$ in $\widetilde{\mathcal{AC}}_n \times (0,\pi]^n \times \mathbb{R}^+$. Suppose first that $0 < \alpha_i < \pi$ for each i. Then each diagonal entry of $B^{(0)}$ is positive except for the bottom right, and after row-reducing $B^{(0)}$ to make it upper-triangular, the bottom-right entry is:

$$2\frac{\cosh J}{\sinh J}\cdot\sum_{i=0}^{n-1}\frac{1-\cos\alpha_i}{\sin\alpha_i}$$

Since each summand is positive, $B^{(0)}$ is non-singular and the claim holds in this case.

If any $\alpha_{j_0}=\pi$ then $\alpha_i<\pi$ for each $i\neq j_0$, since $\alpha_i>0$ for each i and $\sum_{i=0}^{n-1}\alpha_i=2\pi$. In this case the j_0th row of $B^{(0)}$ has all entries equal to 0 but the last, which is $2\sinh J\cosh J>0$. Exchanging the j_0th row of $B^{(0)}$ with the bottom row produces a matrix B' that is almost upper-triangular: only $B'_{ij_0}=-1$ is non-zero among all B'_{ij} with $i<j$. B' is easily row-reduced, along the lines of the previous case, to produce an upper-triangular matrix with non-zero diagonal entries, so the claim holds in this case as well.

In particular, $B^{(0)}$ is invertible at $(d_0,\dots,d_{n-1},f(d_0,\dots,d_{n-1}))$ for each $(d_0,\dots,d_{n-1})$ in $\widetilde{\mathcal{C}}_n\cup\widetilde{\mathcal{BC}}_n$. The implicit function theorem therefore provides a neighborhood V of $(d_0,\dots,d_{n-1})$ in $(\mathbb{R}^+)^n$ and a unique, smooth function $f^{(0)}\colon V\to(0,2\pi)^n\times\mathbb{R}^+$ such that $f^{(0)}(d_0,\dots,d_{n-1})=f(d_0,\dots,d_{n-1})$ and $F\circ(\mathrm{Id}_{\mathbb{R}^n}\times f^{(0)})|_V\equiv(0,\dots,0)$. By uniqueness and (4.1.5), $f^{(0)}=f$ on $V\cap\left(\widetilde{\mathcal{C}}_n\cup\widetilde{\mathcal{BC}}_n\right)$. In particular, since $\widetilde{\mathcal{C}}_n$ is open in $(\mathbb{R}^+)^n$ it follows that $f|_{\widetilde{\mathcal{C}}_n}$ is smooth.

Now define $F^{(1)}\colon\mathbb{R}^n\times(0,2\pi)^n\times\mathbb{R}^+\to\mathbb{R}^{n+1}$ in coordinates by $F^{(1)}=(F^{(1)}_0,F^{(1)}_1,\dots,F^{(1)}_n)$, where $F^{(1)}_i=F^{(0)}_i$ for each $i<n$, and

$$F^{(1)}_n(x_0,\dots,x_{n-1},y_0,\dots,y_{n-1},y_n)=y_0-\sum_{i=1}^{n-1}y_i$$

Since $F^{(1)}$ differs from $F^{(0)}$ only in the final coordinate, for $0\leq i<n$ the identities (4.1.4) still apply to a point $(d_0,\dots,d_{n-1},\alpha_0,\dots,\alpha_{n-1},J)$ taken by $F^{(1)}$ to $(0,\dots,0)\in\mathbb{R}^{n+1}$. On the other hand, $F^{(1)}_n$, takes such a point to 0 if and only if $\alpha_0=\sum_{i=1}^{n-1}\alpha_i$. Therefore as in the case of $F^{(0)}$, Lemma 3.7 and the definition of f imply that:

$$F^{(1)}\circ(\mathrm{Id}_{\mathbb{R}^n}\times f)|_{U_n\cap(\widetilde{\mathcal{AC}}_n-\widetilde{\mathcal{C}}_n)}\equiv(0,\dots,0),\tag{4.1.6}$$

But we note that here we are restricting to a different subset of $\widetilde{\mathcal{AC}}_n$ than in (4.1.5).

At a zero of $F^{(1)}$, $dF^{(1)}=(A^{(0)}\,|\,B^{(1)})$, where $A^{(0)}$ is as above and $B^{(1)}=\left(\frac{\partial F^{(1)}_i}{\partial y_j}\right)^{0\leq i\leq n}_{0\leq j\leq n}$:

$$B^{(1)}=\left(\begin{array}{cccc|c}
\sinh^2 J\cot\left(\frac{\alpha_0}{2}\right) & 0 & \dots & 0 & 2\sinh J\cosh J\\
0 & \sinh^2 J\cot\left(\frac{\alpha_1}{2}\right) & & \vdots & 2\sinh J\cosh J\\
\vdots & & \ddots & 0 & \vdots\\
0 & \dots & 0 & \sinh^2 J\cot\left(\frac{\alpha_{n-1}}{2}\right) & 2\sinh J\cosh J\\
\hline
1 & -1 & \dots & -1 & 0
\end{array}\right)$$

$B^{(1)}$ differs from $B^{(0)}$ only in its bottom left corner (we have simplified it with the identity $\frac{\sin\alpha}{1-\cos\alpha}=\cot(\alpha/2)$). Considerations analogous to those for $F^{(0)}$ reveal that $B^{(1)}$ is invertible if $\alpha_0=\pi$, and if $\alpha_0<\pi$ then it is invertible if and only if $\tan(\alpha_0/2)\neq\sum_{i=1}^{n-1}\tan(\alpha_i/2)$.

Claim 4.1.7. If $\sum_{i=1}^{n-1} \alpha_i = \alpha_0 < \pi$, and $\alpha_i > 0$ for each i, then $\sum_{i=1}^{n-1} \tan(\alpha_i/2) < \tan(\alpha_0/2)$.

Proof of claim. For $0 < \alpha \leq \beta$ with $\alpha + \beta < \pi/2$, an angle-addition identity gives:

$$\tan(\alpha+\beta) = \frac{\tan\alpha + \tan\beta}{1 - \tan\alpha\tan\beta}$$

Since $\alpha < \pi/2 - \beta$ and sin is increasing on $(0, \pi/2)$, we have $\sin\alpha < \sin(\pi/2 - \beta) = \cos\beta$. Similarly, $\cos\alpha > \sin\beta$, and so $0 < \tan\alpha\tan\beta < 1$. Hence $\tan\alpha + \tan\beta < \tan(\alpha+\beta)$, and the claim follows by induction. $\square$

By the claim, $B^{(1)}$ is invertible for as long as $\alpha_0 \leq \pi$. Thus for $(d_0, \dots, d_{n-1}) \in U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$, the implicit function theorem yields a neighborhood V of $(d_0, \dots, d_{n-1})$ in $(\mathbb{R}^+)^n$ and a unique, smooth function $f^{(1)}$ such that $f^{(1)}(d_0, \dots, d_{n-1}) = f(d_0, \dots, d_{n-1})$ and on V:

$$F^{(1)} \circ (\mathrm{Id}_{\mathbb{R}^n} \times f^{(1)}) \equiv (0, \dots, 0)$$

Uniqueness and (4.1.6) imply that $f^{(1)} \equiv f$ on $V \cap U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$. In particular, f is smooth on $U_n \cap (\widetilde{\mathcal{AC}} - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$, since this set is open.

We recall from Corollary 3.5 that $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$ is the disjoint union of components $\sigma^i(U_n) \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$, where σ acts on $\mathbb{R}^n$ by permuting entries: $\sigma(d_0, \dots, d_{n-1}) = (d_1, \dots, d_{n-1}, d_0)$. Of course σ is smooth, and since J is smooth on $U_n \cap (\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$ and $\widetilde{\mathcal{AC}}_n$ and $\widetilde{\mathcal{BC}}_n$ are preserved by σ it follows that $J \circ \sigma^{-i}$ is smooth on $\sigma^i(U_n) \cap (\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$. By Lemma 3.6, $J \colon \widetilde{\mathcal{AC}}_n \to \mathbb{R}$ satisfies $J = J \circ \sigma$, so $J = J \circ \sigma^{-i}$ is smooth on $\sigma^i(U_n) \cap (\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$.

We showed previously that f is smooth on $\widetilde{\mathcal{C}}_n$, so in particular J is smooth here. The paragraph above therefore completes the proof that J is smooth on $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{BC}}_n$. By Lemma 3.7, the functions α_i satisfy $\alpha_i \circ \sigma^j = \alpha_{i-j}$. Since α_i is smooth on $U_n \cap (\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$ for each i, arguing as in the paragraph above establishes that it is smooth on all of $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{BC}}_n$.

For any $(d_0, \dots, d_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$, we produced a neighborhood V of $(d_0, \dots, d_{n-1})$ in $\mathbb{R}^n$ and $f^{(0)}$ on V such that $F^{(0)} \circ (\mathrm{Id}_{\mathbb{R}^n} \times f^{(0)}) \equiv (0, \dots, 0)$ on V and $f^{(0)} = f$ on $V \cap (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n)$. Writing $f^{(0)} = (\alpha_0^{(0)}, \dots, \alpha_{n-1}^{(0)}, J^{(0)})$ in coordinates, the identities (4.1.2) and (4.1.3) hold on V with α_i replaced by $\alpha_i^{(0)}$ and J replaced by $J^{(0)}$. This is because $dF^{(0)} \circ d(\mathrm{Id}_{\mathbb{R}^n} \times f^{(0)})$ is the $(n+1) \times n$ 0-matrix. Setting its j, j entry equal to 0 yields (4.1.3) for $j \in \{0, \dots, n-1\}$, and setting the i, j entry equal to 0, for $i \neq j \in \{0, \dots, n-1\}$, yields (4.1.2).

If $(d_0, \dots, d_{n-1}) \in \widetilde{\mathcal{C}}_n$ then $J \equiv J^{(0)}$ and $\alpha_i \equiv \alpha_i^{(0)}$ for each i on the open neighborhood $V \cap \widetilde{\mathcal{C}}_n$ of $(d_0, \dots, d_{n-1})$, so the derivatives of J and the α_i equal those of $J^{(0)}$ and $\alpha_i^{(0)}$, respectively, at $(d_0, \dots, d_{n-1})$ and in particular satisfy (4.1.2) and (4.1.3).

For $(d_0, \dots, d_{n-1}) \in U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$ and $f^{(1)}$ with $F^{(1)} \circ (\mathrm{Id}_{\mathbb{R}^n} \times f^{(1)}) \equiv (0, \dots, 0)$ on a neighborhood of $(d_0, \dots, d_{n-1})$, if $f^{(1)} = (\alpha_0^{(1)}, \dots, \alpha_{n-1}^{(1)}, J^{(1)})$ then the same identities (4.1.2) and (4.1.3) hold; in this case substituting $\alpha_i^{(1)}$ for α_i and $J^{(1)}$ for J. This is because the first n coordinates of $F^{(0)}$ and $F^{(1)}$ agree. It follows as above that if $(d_0, \dots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n)$ then the derivatives of $J = J^{(1)}$ and $\alpha_i = \alpha_i^{(1)}$ also satisfy (4.1.2) and (4.1.3) here.

Around $(d_0,\dots,d_{n-1})\in U_n\cap\widetilde{BC}_n=\mathrm{Graph}(b_0)$ there is a single neighborhood V and smooth functions $f^{(0)}$ and $f^{(1)}$ on V such that $F^{(\epsilon)}\circ(\mathrm{Id}_{\mathbb{R}^n}\times f^{(\epsilon)})\equiv(0,\dots,0)$ on V for $\epsilon\in\{0,1\}$, and $f=f^{(0)}$ on $V\cap(\widetilde{\mathcal{C}}_n\cup\widetilde{\mathcal{BC}}_n)$ and $f=f^{(1)}$ on $V\cap(\widetilde{\mathcal{AC}}_n-\widetilde{\mathcal{C}}_n)$. In particular,

$$f^{(0)}(d_0,\dots,d_{n-1})=f(d_0,\dots,d_{n-1})=f^{(1)}(d_0,\dots,d_{n-1}),$$

so at least f is continuous here. In fact we will show that at such a point the identities (4.1.2) and (4.1.3) are enough to prescribe that J is $C^{(1)}$ here, as are the α_i for $i>0$.

For $(d_0,\dots,d_{n-1})\in U_n\cap\mathcal{BC}_n=\mathrm{Graph}(b_0)$, Lemma 3.9 gives $\alpha_0(d_0,\dots,d_{n-1})=\pi$ and $J(d_0,\dots,d_{n-1})=d_0/2$. Therefore for $j>0$, taking $i=0$ in the identity (4.1.2) yields:

$$\frac{\partial J^{(0)}}{\partial d_j}(d_0,\dots,d_{n-1})=\frac{\partial J^{(1)}}{\partial d_j}(d_0,\dots,d_{n-1})=0$$

Here we are using the fact that (4.1.2) applies to the functions $f^{(0)}$ and $f^{(1)}$ at $(d_0,\dots,d_{n-1})$, with appropriate substitutions of indices. Similarly, taking $j=0$ in (4.1.3) gives:

$$\frac{\partial J^{(0)}}{\partial d_0}(d_0,\dots,d_{n-1})=\frac{\partial J^{(1)}}{\partial d_0}(d_0,\dots,d_{n-1})=\frac{1}{2}$$

To conclude that J is differentiable here we use the following elementary calculus fact:

Fact 4.2. If $f\colon\mathbb{R}\to\mathbb{R}$ is continuous at x and has well-defined derivatives from the left and the right here, and these are equal, then f is differentiable at x.

Applying this to the composition of J with $\iota_0\colon t\mapsto(d_0+t,d_1,\dots,d_{n-1})$ shows that $\partial J/\partial d_0$ is well-defined and equal to $1/2$, since it is not hard to see that $J\circ\iota_0=J^{(0)}\circ\iota_0$ for $t\le 0$ and $J\circ\iota_0=J^{(1)}\circ\iota_0$ for $t\ge 0$. Defining ι_j analogously for $j>0$, we find that $J\circ\iota_j=J^{(1)}\circ\iota_j$ for $t\le 0$ and $J\circ\iota_j=J^{(0)}\circ\iota_j$ for $t\ge 0$; thus $\partial J/\partial d_j$ exists here and is zero.

It follows that for each j, $\partial J/\partial d_j$ is continuous at $(d_0,\dots,d_{n-1})$, since it agrees with $\partial J^{(0)}/\partial d_j$ on $V\cap\widetilde{\mathcal{C}}_n$, with $\partial J^{(1)}/\partial d_j$ on $V\cap(\widetilde{\mathcal{AC}}_n-(\widetilde{\mathcal{C}}_n\cup\widetilde{\mathcal{BC}}_n))$, and with both on $V\cap\widetilde{\mathcal{BC}}_n$ by above.

Again taking $(d_0,\dots,d_{n-1})\in\mathrm{Graph}(b_0)$, fixing $i>0$ and inserting the values of $\partial J^{(\epsilon)}/\partial d_j$ from above, $\epsilon\in\{0,1\}$, into the inequalities (4.1.2) and (4.1.3) yields:

$$\frac{\partial\alpha_i^{(\epsilon)}}{\partial d_0}=-\coth(d_0/2)\tan(\alpha_i/2)\qquad\frac{\partial\alpha_i^{(\epsilon)}}{\partial d_j}=0\quad(j\neq 0,i)\qquad\frac{\partial\alpha_i^{(\epsilon)}}{\partial d_i}=\coth(d_i/2)\tan(\alpha_i/2)$$

Since α_i agrees with one or the other of $\alpha_i^{(\epsilon)}$ on a neighborhood of $(d_0,\dots,d_{n-1})$, we may argue as with J to show that α_i is C^1 here and its derivatives are given above.

The paragraphs above have determined the values of $\partial J/\partial d_j$ and $\partial\alpha_i/\partial d_j$ on $\mathrm{Graph}(b_0)$, for any $j\in\{0,\dots,n-1\}$ and any such i that is also greater than 0. We used each of the n^2 identities (4.1.2) and (4.1.3) to produce these descriptions, and in particular they contain no further information that might aid in obtaining the partials of α_0. In confirming the identity (4.1.1), we will find that these are not well-defined here.

For $(d_0,\dots,d_{n-1})\in\overline{U}_n\cap(\widetilde{\mathcal{C}}_n\cup\widetilde{\mathcal{BC}}_n)$, let $f^{(0)}=(\alpha_0^{(0)},\dots,\alpha_{n-1}^{(0)},J^{(0)})$ agree with f at $(d_0,\dots,d_{n-1})$ and satisfy $F^{(0)}\circ(\mathrm{Id}_{\mathbb{R}^n}\times f^{(0)})=(0,\dots,0)$ on a neighborhood V. For each

$j \in \{0, \ldots, n-1\}$, setting the $(n-1, j)$ entry of $d(F^{(0)} \circ (\mathrm{Id}_{\mathbb{R}^n} \times f^{(0)}))$ equal to 0 yields:

$$0 = \sum_{k=0}^{n-1} \frac{\partial \alpha_k^{(0)}}{\partial d_j}(d_0, \ldots, d_{n-1})$$

If $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{C}}_n$ then since $f = f^{(0)}$ on $V \cap \widetilde{\mathcal{C}}_n$ they have the same derivative here. In particular, the $\partial\alpha_k/\partial d_j$ satisfy the identity above at $(d_0, \ldots, d_{n-1})$. Solving this for $\partial\alpha_0/\partial d_j$ confirms the relevant case of (4.1.1). If $(d_0, \ldots, d_{n-1}) \in U_n \cap \widetilde{\mathcal{BC}}_n = \mathrm{Graph}(b_0)$, then since $\alpha_0 = \alpha_0^{(0)}$ on $V \cap (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n)$, computing $\partial\alpha_0/\partial d_j$ from within this set gives a well-defined value equal to that of $\partial\alpha_0^{(0)}/\partial d_j$ here.

For $(d_0, \ldots, d_{n-1}) \in \overline{U}_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$, let $f^{(1)} = (\alpha_0^{(1)}, \ldots, \alpha_{n-1}^{(1)}, J^{(1)})$ satisfy $f^{(1)} = f$ at $(d_0, \ldots, d_{n-1})$ and $F^{(1)} \circ (\mathrm{Id}_{\mathbb{R}^n} \times f^{(1)}) \equiv (0, \ldots, 0)$ on a neighborhood of it. Then for any $j \in \{0, \ldots, n-1\}$, setting the $(n-1, j)$ entry of $d(F^{(1)} \circ (\mathrm{Id}_{\mathbb{R}^n} \times f^{(1)}))$ equal to 0 gives:

$$0 = \frac{\partial \alpha_0^{(1)}}{\partial d_j}(d_0, \ldots, d_{n-1}) - \sum_{k=1}^{n-1} \frac{\partial \alpha_k^{(1)}}{\partial d_j}(d_0, \ldots, d_{n-1})$$

If $(d_0, \ldots, d_{n-1}) \in U_n \cap (\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$ then $f = f^{(1)}$ on a neighborhood of $(d_0, \ldots, d_{n-1})$, so here $df = df^{(1)}$ and the relevant case of (4.1.1) is confirmed. At $(d_0, \ldots, d_{n-1}) \in \mathrm{Graph}(b_0)$, computing $\partial\alpha_0/\partial d_j$ from within $V \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$ yields $\partial\alpha_0^{(1)}/\partial d_j$ for each j.

At $(d_0, \ldots, d_{n-1}) \in \mathrm{Graph}(b_0)$, we have already showed that $\partial\alpha_k^{(0)}/\partial d_j = \partial\alpha_k^{(1)}/\partial d_j$ for each j, as long as $k > 0$. The two identities above thus imply that $\partial\alpha_0^{(1)}/\partial d_j = -\partial\alpha_0^{(0)}/\partial d_j$ here for each j. This proves all claims of the proposition on $\overline{U}_n$.

We will extend the proposition to $\sigma^{-i}(\overline{U}_n)$ for arbitrary $i \in \{0, \ldots, n-1\}$ using the fact that $\alpha_k = \alpha_{k-i} \circ \sigma^i$. Recall that by definition $\sigma^{-i}(d_0, \ldots, d_{n-1}) = (d_{n-i}, \ldots, d_{n-1}, d_0, \ldots, d_{n-1-i})$. In particular, on $\sigma^{-i}(\mathrm{Graph}(b_0))$ we have $\alpha_i \equiv \pi$. Also, $\frac{\partial(\sigma^i)_k}{\partial d_j} = \delta_{j-i}^k$, so by the chain rule,

$$\begin{aligned}
\frac{\partial \alpha_k}{\partial d_j}(d_0, \ldots, d_{n-1}) &= \sum_{m=0}^{n-1} \frac{\partial \alpha_{k-i}}{\partial d_m}\sigma^i(d_0, \ldots, d_{n-1}) \cdot \frac{\partial(\sigma^i)_m}{\partial d_j}(d_0, \ldots, d_{n-1}) \\
&= \frac{\partial \alpha_{k-i}}{\partial d_{j-i}}\sigma^i(d_0, \ldots, d_{n-1})
\end{aligned}$$

In particular, at $(d_0, \ldots, d_{n-1}) \in \sigma^{-i}(\overline{U}_n) - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n)$, if $\sigma^i(d_0, \ldots, d_{n-1}) = (d'_0, \ldots, d'_{n-1}) \in \overline{U}_n$, then for any $j \in \{0, \ldots, n-1\}$:

$$\begin{aligned}
\frac{\partial \alpha_i}{\partial d_j}(d_0, \ldots, d_{n-1}) &= \frac{\partial \alpha_0}{\partial d_{j-i}}(d'_0, \ldots, d'_{n-1}) \\
&= \begin{cases} -\sum_{k\neq 0} \frac{\partial \alpha_k}{\partial d_{j-i}}(d'_0, \ldots, d'_{n-1}) & (d'_0, \ldots, d'_{n-1}) \in \overline{U}_n \cap \widetilde{\mathcal{C}}_n \\ \sum_{k\neq 0} \frac{\partial \alpha_k}{\partial d_{j-i}}(d'_0, \ldots, d'_{n-1}) & (d'_0, \ldots, d'_{n-1}) \in \overline{U}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n) \end{cases} \\
&= \begin{cases} -\sum_{k\neq i} \frac{\partial \alpha_k}{\partial d_j}(d_0, \ldots, d_{n-1}) & (d_0, \ldots, d_{n-1}) \in \sigma^{-i}(\overline{U}_n) \cap \widetilde{\mathcal{C}}_n \\ \sum_{k\neq i} \frac{\partial \alpha_k}{\partial d_j}(d_0, \ldots, d_{n-1}) & (d_0, \ldots, d_{n-1}) \in \sigma^{-i}(\overline{U}_n) - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n) \end{cases}
\end{aligned}$$

Above we have used the chain rule twice and the $\overline{U}_n$ case of (4.1.1) once. This confirms the remaining cases of (4.1.1), and hence the proposition. $\square$

Corollary 4.3. *For $n \geq 3$ and $i \in \{0, \dots, n-1\}$, the function β_i from Lemma 3.7 is smooth on $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{BC}}_n$, C^0 on $\widetilde{\mathcal{AC}}_n$, and C^1 on $\widetilde{\mathcal{AC}}_n - \sigma^{-i}(\mathrm{Graph}(b_0))$. Its derivatives satisfy:*

$$\frac{\partial \beta_i}{\partial d_i} = \frac{-1}{2\cosh J}\frac{\partial \alpha_i}{\partial d_i} - \frac{1}{2}\sqrt{\frac{1}{\cosh^2(d_i/2)} - \frac{1}{\cosh^2 J}} \qquad \frac{\partial \beta_i}{\partial d_j} = \frac{-1}{2\cosh J}\frac{\partial \alpha_i}{\partial d_j} \quad (j \neq i)$$

At $(d_0, \dots, d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$, $\frac{\partial \beta_i}{\partial d_j}$ has well-defined values computed from within $\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$ or $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$, and these are opposite.

Proof. Rearranging the relation between α_i and β_i from Lemma 3.7 yields:

$$\beta_i \equiv \sin^{-1}\left(\frac{\sin\alpha_i \sinh J}{\sinh d_i}\right) \tag{4.3.1}$$

Recall that α_i, d_i and J satisfy the identity (4.1.4), which implies:

$$\frac{\sin\alpha_i \sinh J}{\sinh d_i} = \frac{\sin\alpha_i}{\sinh d_i}\sqrt{\frac{\cosh d_i - 1}{1 - \cos\alpha_i}} = \sqrt{\frac{1+\cos\alpha_i}{1+\cosh d_i}} \in [0,1)$$

Since the inverse sine is smooth on $[0,1)$, appealing to (4.3.1) we find that the smoothness and continuity properties of β_i depend on those of α_i and J. Hence β_i is continuous on $\widetilde{\mathcal{AC}}_n$, smooth on $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{BC}}_n$, and C^1 on $\widetilde{\mathcal{AC}}_n - \sigma^{-i}(\mathrm{Graph}(b_0))$.

In order to relate the derivatives of β_i to those of α_i, we use the identity (4.3.1), together with the chain and product rules. For $j \neq i$, after some simplification this yields:

$$\frac{\partial \beta_i}{\partial d_j} = \frac{\sinh d_i}{\sqrt{\sinh^2 d_i - \sin^2\alpha_i \sinh^2 J}}\left[\frac{\sinh J\cos\alpha_i}{\sinh d_i}\frac{\partial \alpha_i}{\partial d_j} + \frac{\sin\alpha_i\cosh J}{\sinh d_i}\frac{\partial J}{\partial d_j}\right] \tag{4.3.2}$$

$$= \frac{\sinh J\sin\alpha_i}{\sqrt{\sinh^2 d_i - \sin^2\alpha_i \sinh^2 J}}\left[\cot\alpha_i\frac{\partial \alpha_i}{\partial d_j} + \coth J\frac{\partial J}{\partial d_j}\right] \tag{4.3.3}$$

We simplify using the identity (4.1.4) from the proof of Proposition 4.1. Substituting for instances of $\sinh J$ and simplifying gives:

$$\frac{\partial \beta_i}{\partial d_j} = \sqrt{\frac{1+\cos\alpha_i}{\cosh d_i - \cos\alpha_i}}\left[\cot\alpha_i\frac{\partial \alpha_i}{\partial d_j} + \coth J\frac{\partial J}{\partial d_j}\right]$$

In order to further simplify we will use the trigonometric identity below:

$$\cot\alpha = \frac{1}{2}\cot(\alpha/2) - \frac{1-\cos\alpha}{2\sin\alpha} = \frac{1}{2}\cot(\alpha/2) - \frac{1}{2}\sqrt{\frac{1-\cos\alpha}{1+\cos\alpha}}$$

This may be somewhat non-standard but follows easily from the standard identities. Inserting it into the expression above allows the application of (4.1.2), after which we have:

$$\frac{\partial \beta_i}{\partial d_j} = \frac{-1}{2}\sqrt{\frac{1-\cos\alpha_i}{\cosh d_i - \cos\alpha_i}}\frac{\partial \alpha_i}{\partial d_j} = \frac{-1}{2\cosh J}\frac{\partial \alpha_i}{\partial d_j}$$

The latter equality above follows again from (4.1.4).

Obtaining the identity for $j = i$ follows the same chain of argument as in the case above, with two differences. First, the product rule gives an additional summand in (4.3.2), since $\partial d_i/\partial d_i = 1$. After factoring, this yields the additional term $-\cosh d_i/\sinh d_i$ inside the brackets in (4.3.3). Second, we must apply (4.1.3) instead of (4.1.2). The result:

$$\frac{\partial \beta_i}{\partial d_i} = \frac{-1}{2\cosh J}\frac{\partial \alpha_i}{\partial d_i} + \sqrt{\frac{1+\cos\alpha_i}{\cosh d_i - \cos\alpha_i}}\left[\frac{1}{2}\coth(d_i/2) - \frac{\cosh d_i}{\sinh d_i}\right]$$

Upon applying "double angle" identities, the expression in brackets above simplifies to $-\sinh(d_i/2)/2\cosh(d_i/2)$. Solving the identity (4.1.4) for $\cos\alpha_i$ and substituting above, and replacing the remaining instance of $\cosh d_i$ with $2\sinh^2(d_i/2)+1$ yields:

$$\frac{\partial \beta_i}{\partial d_i} = \frac{-1}{2\cosh J}\frac{\partial \alpha_i}{\partial d_i} - \frac{1}{2}\sqrt{\frac{1}{\cosh^2(d_i/2)} - \frac{1}{\cosh^2 J}}$$

At $(d_0, \dots, d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$, Proposition 4.1 implies that α_i has well-defined partials computed from within $\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$ or $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$; call them $\partial\alpha_i^-/\partial d_j$ and $\partial\alpha_i^+/\partial d_j$, respectively. The same therefore holds for β_i, and by the above we have:

$$\frac{\partial \beta_i^-}{\partial d_j} = \frac{-1}{2\cosh J}\frac{\partial \alpha_i^-}{\partial d_j} = \frac{1}{2\cosh J}\frac{\partial \alpha_i^+}{\partial d_j} = -\frac{\partial \beta_i^+}{\partial d_j}$$

Here we have used the assertion of Proposition 4.1, that $\partial\alpha_i^-/\partial d_j = -\partial\alpha_i^+/\partial d_j$ for each j. For $j = i$ we have also used Lemma 3.9, which asserts that $J(d_0, \dots, d_{n-1}) = d_i/2$ for $(d_0, \dots, d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$, so here $\cosh J = \cosh(d_i/2)$. □

Proposition 4.4. *For $n \geq 3$, ν_i is smooth on $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{BC}}_n$ and C^1 on $\widetilde{\mathcal{AC}}_n$.*

Proof. The definition in Lemma 3.8 implies that $\nu_i \equiv \beta_i + \beta_{i+1}$ on $\widetilde{\mathcal{C}}_n$, so it is smooth here by Corollary 4.3. That ν_i is smooth on $U_n \cap (\widetilde{\mathcal{AC}} - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$ for each i also follows directly from the definitions and Corollary 4.3. The identity $\nu_i = \nu_{i-j} \circ \sigma^j$ implies that smoothness holds on all of $\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n)$.

For $i \in \{1, \dots, n-2\}$, ν_i is continuously differentiable at $(d_0, \dots, d_{n-1}) \in \mathrm{Graph}(b_0)$ since β_i and β_{i+1} are. Since β_0 is not differentiable here, the conclusion is not immediate for ν_0 or ν_{n-1}. However, Lemma 3.9 implies that $\beta_0 = 0$, and hence ν_0 and ν_{n-1} are continuous, here.

Furthermore, Corollary 4.3 implies that β_0 has well-defined partial derivatives $\partial\beta_0^-/\partial d_j$ and $\partial\beta_0^+/\partial d_j$ at $(d_0, \dots, d_{n-1}) \in \mathrm{Graph}(b_0)$ for each j, computed from within $\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$ and $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$, respectively; and that $\partial\beta_0^-/\partial d_j = -\partial\beta_0^+/\partial d_j$ here. This implies:

$$\left(\frac{\partial \beta_1}{\partial d_j} + \frac{\partial \beta_0^-}{\partial d_j}\right)(d_0, \dots, d_{n-1}) = \left(\frac{\partial \beta_1}{\partial d_j} - \frac{\partial \beta_0^+}{\partial d_j}\right)(d_0, \dots, d_{n-1})$$

Using Fact 4.2 as in the proof of Proposition 4.1, we thus find that ν_0 is continuously differentiable here. An analogous argument establishes that ν_{n-1} is as well. The $\mathbb{Z}_n$-equivariance property of the ν_i may now be applied to show that they are C^1. □

Using the identities of Proposition 4.1, one may produce an explicit formula for the partial derivatives of J. This in turn gives rise to the estimates of the lemma below.

Lemma 4.5. *For $n \geq 3$ and $(d_0, \ldots, d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$,*

$$\frac{\partial J}{\partial d_j}(d_0, \ldots, d_{n-1}) = \begin{cases} \frac{1}{2} & j = i \\ 0 & j \neq i \end{cases}$$

(Here $J\colon \widetilde{\mathcal{AC}}_n \to \mathbb{R}^+$ is from Lemma 3.6.) At any $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{BC}}_n$, if $d_i > d_j$ then $|\partial J/\partial d_i| > |\partial J/\partial d_j|$. Furthermore, on $\widetilde{\mathcal{C}}_n$, $0 < \partial J/\partial d_j < \frac{1}{2}$ for each j; and on $\sigma^{-i}(U_n) \cap (\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$, $\partial J/\partial d_i > \frac{1}{2}$ and $\partial J/\partial d_j < 0$ for $j \neq i$.

Proof. Lemma 3.9 implies that $\sigma^{-i}(\mathrm{Graph}(b_0))$ is the zero-locus of $\cot(\alpha_i/2)$. Here the identities (4.1.2) thus give $\partial J/\partial d_j = 0$ for $j \neq i$, and (4.1.3) gives:

$$\frac{\partial J}{\partial d_i} = \frac{\coth(d_j/2)}{2 \coth J} = \frac{1}{2}$$

The latter equality above follows again from Lemma 3.9, which asserts that $J \equiv d_i/2$ on $\sigma^{-i}(\mathrm{Graph}(b_0))$. Since $\widetilde{\mathcal{BC}}_n = \bigcup_{i=0}^{n-1} \sigma^{-i}(\mathrm{Graph}(b_0))$ by definition, we have produced a complete description of the values of $\partial J/\partial d_j$ there. We will assume below that $(d_0, \ldots, d_{n-1}) \notin \widetilde{\mathcal{BC}}_n$, so in particular $\cot(\alpha_i/2) \neq 0$ at $(d_0, \ldots, d_{n-1})$ for each i, by Lemma 3.9.

For any fixed j, solving the identities (4.1.2) and (4.1.3) for $\partial \alpha_i/\partial d_j$ gives:

$$\frac{\partial \alpha_i}{\partial d_j} = \begin{cases} -2\tan(\alpha_i/2)\coth J \frac{\partial J}{\partial d_j} & i \neq j \\ \tan(\alpha_j/2)\left(\coth(d_j/2) - 2\coth J \frac{\partial J}{\partial d_j}\right) & i = j \end{cases} \tag{4.5.1}$$

For $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{C}}_n$, inserting the formulas above into the relevant case of (4.1.1) and solving for J gives:

$$\frac{\partial J}{\partial d_j}(d_0, \ldots, d_{n-1}) = \frac{\coth(d_j/2)\tanh J \tan(\alpha_j/2)}{2\sum_{i=0}^{n-1} \tan(\alpha_i/2)}$$

Since all terms of this formula are positive for each j, we conclude that $\partial J/\partial d_j > 0$. Recall that for any j, J, d_j, and α_j satisfy the identity (4.1.4), which implies that if $d_i > d_j$ then $\alpha_i > \alpha_j$, and also (using half-angle identities) that $\sinh(d_j/2) = \sinh J \sin(\alpha_j/2)$. Thus:

$$\coth(d_j/2)\tanh J \tan(\alpha_j/2) = \frac{\cosh(d_j/2)}{\cosh J \cos(\alpha_j/2)}$$

Substituting this into the formula for $\partial J/\partial d_j$ gives:

$$\frac{\partial J}{\partial d_j} = \frac{\cosh(d_j/2)\,\mathrm{sech}\, J}{2\cos(\alpha_j/2)\sum_{i=0}^{n-1}\tan(\alpha_i/2)} = \frac{\cosh(d_j/2)\,\mathrm{sech}\, J}{2\left[\sin(\alpha_j/2) + \cos(\alpha_j/2)\sum_{i\neq j}\tan(\alpha_i/2)\right]}$$

Since $0 < \alpha_i < \pi$ for each i, we have $\tan(\alpha_i/2) > \sin(\alpha_i/2)$. Since $\alpha_j < \pi$ and $\sum_{i=0}^{n-1} \alpha_i = 2\pi$, we have $\sum_{i\neq j} \alpha_i > \pi$, so Claim 2.3.1 implies that $\sum_{i \neq j} \sin(\alpha_i/2) > 1$. Putting these together, we find:

$$\sin(\alpha_j/2) + \cos(\alpha_j/2)\sum_{i\neq j}\tan(\alpha_i/2) > \sin(\alpha_j/2) + \cos(\alpha_j/2) > 1$$

Inserting into the expression above gives $\partial J/\partial d_j < \frac{1}{2}\cosh(d_j/2)\,\mathrm{sech}\, J < 1/2$ on $\widetilde{\mathcal{C}}_n$.

Now suppose $(d_0, \dots, d_{n-1}) \in U_n \cap (\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$. For fixed j, inserting the data from (4.5.1) into the relevant case of (4.1.1) and solving for $\partial J/\partial d_j$ yields:

$$\frac{\partial J}{\partial d_0} = \frac{\coth(d_0/2)\tanh J \tan(\alpha_0/2)}{2\left[\tan(\alpha_0/2) - \sum_{i=1}^{n-1}\tan(\alpha_i/2)\right]} \quad \frac{\partial J}{\partial d_j} = \frac{-\coth(d_j/2)\tanh J \tan(\alpha_j/2)}{2\left[\tan(\alpha_0/2) - \sum_{i=1}^{n-1}\tan(\alpha_i/2)\right]} \ (j > 0)$$

Since Lemma 3.7 asserts in particular that $\alpha_0 \equiv \sum_{i=1}^{n-1} \alpha_i$ on $U_n \cap (\widetilde{\mathcal{AC}}_n - \mathcal{C}_n)$, Claim 4.1.7 implies that $\tan(\alpha_0/2) > \sum_{i=1}^{n-1} \tan(\alpha_i/2)$ on $U_n \cap (\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$. Thus $\partial J/\partial d_j < 0$ here for $j > 0$. Lemma 3.9 implies that $J > d_0/2$ outside $\widetilde{\mathcal{BC}}_n$, so since the hyperbolic tangent is increasing, $\coth(d_0/2)\tanh J > 1$ and $\partial J/\partial d_0 > 1/2$ on $U_n \cap (\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$.

For $i > 0$ and $(d_0, \dots, d_{n-1}) \in \sigma^{-i}(U_n) \cap (\widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n))$, we use the fact that $J = J \circ \sigma^{-i}$ here, where $\sigma^{-i}(d_0, \dots, d_{n-1}) = (d_{n-i}, \dots, d_{n-1}, d_0, \dots, d_{n-i-1})$. Together with the chain rule, this shows that if $(d'_0, \dots, d'_{n-1}) = \sigma^i(d_0, \dots, d_{n-1})$ then:

$$\frac{\partial J}{\partial d_j}(d_0, \dots, d_{n-1}) = \frac{\partial J}{\partial d_{j-i}}(d'_0, \dots, d'_{n-1})$$

Applying the U_n case above, since the α_j satisfy $\alpha_k = \alpha_{k+i} \circ \sigma^{-i}$, we obtain an analogous description of $\partial J/\partial d_j$. This implies the desired bounds.

Finally, at any $(d_0, \dots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{BC}}_n$, applying the descriptions above yields:

$$\frac{|\partial J/\partial d_i|}{|\partial J/\partial d_j|} = \frac{\cosh(d_i/2)\cos(\alpha_j/2)}{\cosh(d_j/2)\cos(\alpha_i/2)}$$

If $d_i > d_j$, then $\alpha_i > \alpha_j$, and since cos is decreasing we have $|\partial J/\partial d_i| > |\partial J/\partial d_j|$. $\square$

Corollary 4.6. *For $n \geq 3$, $(d_0, \dots, d_{n-1}) \in U_n \cap \widetilde{\mathcal{AC}}_n$ is in $\widetilde{\mathcal{C}}_n$ if and only if $J(d_0, \dots, d_{n-1}) < \frac{1}{2}b_0(d_1, \dots, d_{n-1})$, for b_0 as in Lemma 3.4.*

Proof. For fixed $(d_1, \dots, d_{n-1}) \in (\mathbb{R}^+)^{n-1}$, Lemma 4.5 implies that $d_0 \mapsto J(d_0, d_1, \dots, d_{n-1})$ is increasing. Lemma 3.4 implies that $J(B_0, d_1, \dots, d_{n-1}) = B_0/2$, where $B_0 = b_0(d_1, \dots, d_{n-1})$, so $J(d_0, d_1, \dots, d_{n-1}) < B_0/2$ if and only if $d_0 < B_0$. The conclusion now follows from Corollary 3.11. $\square$

Lemma 4.7. *For $n \geq 3$, $\widetilde{\mathcal{HC}}_n$ has a neighborhood V in $\mathbb{R}^n$ with $V \cap \widetilde{\mathcal{AC}}_n \subset \widetilde{\mathcal{AC}}_n - (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n)$; and $J(d_0, \dots, d_{n-1}) \to \infty$ as $(d_0, d_1, \dots d_{n-1})$ in $\widetilde{\mathcal{AC}}_n$ approaches $\widetilde{\mathcal{HC}}_n$.*

Proof. Fix $(H_0, d_0^{(0)}, \dots, d_{n-1}^{(0)})$ satisfying $d_i > 0$ for all i and $\sinh(H_0/2) = \sum_{i=1}^{n-1} \sinh(d_i/2)$. Lemma 3.10 implies that $(H_0, d_0^{(0)}, \dots, d_{n-1}^{(0)}) \in \mathrm{Graph}(h_0) \subset \widetilde{\mathcal{HC}}_n$, and since $H_0 > d_i$ for each i, Lemma 2.3 applies and implies that $\sum_{i=1}^{n-1} A_{d_i^{(0)}}(H_0/2) < \pi$.

By Lemma 2.1, the map $(d, J) \mapsto A_d(J)$ is continuous on $\{(d, J) \in \mathbb{R}^2 \mid J \geq d/2\}$. It follows that $(H_0, d_1^{(0)}, \dots, d_{n-1}^{(0)})$ has a neighborhood in $\mathbb{R}^n$ on which

$$(d_0, \dots, d_{n-1}) \mapsto \sum_{i=1}^{n-1} A_{d_i}(d_0/2)$$

is well-defined and continuous; and therefore that a smaller neighborhood $V(H_0, d_1, \ldots, d_{n-1})$ is contained in the preimage of $(0, \pi)$ under this map. The union over Graph(h_0) of such neighborhoods is a neighborhood V_0 of Graph(h_0) with $V_0 \cap \widetilde{\mathcal{AC}}_n \subset \widetilde{\mathcal{AC}}_n - \overline{\mathcal{C}}_n$ by Lemma 3.3. Since the $\mathbb{Z}_n$-action preserves $\widetilde{\mathcal{AC}}_n$ and $\widetilde{\mathcal{C}}_n$, $V = \mathbb{Z}_n \cdot V_0$ satisfies the conclusion above.

Lemma 2.1 asserts that:

$$\lim_{J\to\infty} \frac{\sum_{i=1}^{n-1} A_{d_i^{(0)}}(J)}{A_{H_0}(J)} = \lim_{J\to\infty} \frac{\sum_{i=1}^{n-1} A'_{d_i^{(0)}}(J)}{A'_{H_0}(J)} = \frac{\sum_{i=1}^{n-1} \sinh(d_i^{(0)}/2)}{\sinh(H_0/2)} = 1,$$

and also that the ratio of derivatives above is strictly increasing. Thus $|\sum_{i=1}^{n-1} A'_{d_i^{(0)}}(J)| < |A'_{H_0}(J)|$ at each $J \in [H_0/2, \infty)$, so A_{H_0} decreases faster than $\sum_{i=1}^{n-1} A_{d_i^{(0)}}$ on all of $[H_0/2, \infty)$. If A_{H_0} took the same value as $\sum_{i=1}^{n-1} A_{d_i^{(0)}}$ at some J in this interval, then the limit of their ratios would be less than 1, contradicting the above. Therefore $A_{H_0} - \sum_{i=1}^{n-1} A_{d_i^{(0)}}$ is a positive, decreasing function of J on $[H_0/2, \infty)$ with a horizontal asymptote of 0.

Continuity of $(d, J) \mapsto A_d(J)$ implies that as $(d_0, d_1, \ldots d_{n-1}) \to (H_0, d_1^{(0)}, \ldots, d_{n-1}^{(0)})$, the functions A_{d_0} and A_{d_i} converge to A_{H_0} and $A_{d_i^{(0)}}$, respectively, uniformly on compact subsets. More precisely, for any $\epsilon > 0$ and d_0 with $|d_0 - H_0| < 2\epsilon$, the function A_{d_0} is defined on $[H_0/2 + \epsilon, \infty)$ and converges to A_{H_0} on compact subsets of this interval as $d_0 \to H_0$. By the above, for ϵ small enough and $|(d_0, d_1, \ldots, d_{n-1}) - (H_0, d_1^{(0)}, \ldots, d_{n-1}^{(0)})| < 2\epsilon$ we may assume that $(d_0, \ldots, d_{n-1}) \in \mathcal{AC}_n - \overline{\mathcal{C}}_n$ for all k, and hence that $d_i < d_0$ for each $i > 0$. Then A_{d_i} is also defined on $[H_0/2 + \epsilon, \infty)$ and converges to $A_{d_i^{(0)}}$ uniformly on compact subsets thereof.

For $J_0 > H_0/2$, let $B_0 = A_{H_0}(J_0) - \sum_{i=1}^{n-1} A_{d_i^{(0)}}(J_0)$, so that B_0 is a lower bound for the values of $A_{H_0} - \sum_{i=1}^{n-1} A_{d_i^{(0)}}$ on $[H_0/2, J_0]$. Then by the above there is an $\epsilon > 0$ such that if $|(d_0, d_1, \ldots, d_{n-1}) - (H_0, d_1^{(0)}, \ldots, d_{n-1}^{(0)})| < 2\epsilon$ then $A_{d_0} - \sum_{i=1}^{n-1} A_{d_i}$ is defined on $[H_0/2 + \epsilon, \infty)$ and within $B_0/2$ of $A_{H_0} - \sum_{i=1}^{n-1} A_{d_i}$ on $[H_0/2 + \epsilon, J_0]$. In particular, it is positive here. By Lemma 1.10 the radius J_1 of $(d_0, \ldots, d_{n-1})$ satisfies $A_{d_0}(J_1) = \sum_{i=1}^{n-1} A_{d_i}(J_1)$, and by Lemma 2.2 $A_{d_0}(J) < \sum_{i=1}^{n-1} A_{d_i}(J)$ for all $J > J_1$. It follows that $J_1 > J_0$ for all such $(d_0, \ldots, d_{n-1})$. By its definition in Lemma 3.6, $J(d_0, \ldots, d_{n-1}) = J_1$, and the lemma follows. □

Lemma 4.8. *For $n \geq 3$, the functions α_i and β_i on $\widetilde{\mathcal{AC}}_n$ extend continuously to $\widetilde{\mathcal{HC}}_n$ by:*

$$\begin{aligned} &\alpha_i(d_0, \ldots, d_{n-1}) = 0 \\ &\beta_i(d_0, \ldots, d_{n-1}) = \sin^{-1}\left(\frac{1}{\cosh(d_i/2)}\right) \qquad\qquad (d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{HC}}_n \end{aligned}$$

Furthermore, their extensions satisfy $\alpha_i = \alpha_{i-j} \circ \sigma^j$ and $\beta_i = \beta_{i-j} \circ \sigma^j$.

Proof. Fix $(d_1^{(0)}, \ldots, d_{n-1}^{(0)}) \in (\mathbb{R}^+)^{n-1}$ and let $H_0 = h_0(d_1^{(0)}, \ldots, d_{n-1}^{(0)})$. Recall from the proof of Proposition 4.1 that α_i satisfies (4.1.4) on $\widetilde{\mathcal{AC}}_n$. Applying half-angle identities gives:

$$\sin(\alpha_i/2) \sinh J = \sinh(d_i/2)$$

For any fixed i, as $(d_0, d_1, \dots d_{n-1}) \to (H_0, d_1^{(0)}, \dots, d_{n-1}^{(0)})$ the right-hand side of this equation remains bounded. Since Lemma 4.7 implies $J \to \infty$, it follows that $\sin(\alpha_i/2)$, and hence also α_i, approaches 0.

We may rewrite the identity from Lemma 3.7 that $\beta_i(d_0, \dots, d_{n-1})$ satisfies as:

$$\sin\beta_i \sinh d_i = 2\sin(\alpha_i/2)\sinh J\cos(\alpha_i/2) = 2\sinh(d_i/2)\cos(\alpha_i/2)$$

The first equation above uses the double angle identity for sine, and the second uses the revised (4.1.4) above. As $(d_0, d_1, \dots d_{n-1}) \to (H_0, d_1^{(0)}, \dots, d_{n-1}^{(0)})$, $\cos(\alpha_i/2) \to 1$ since $\alpha_i \to 0$. The "double angle identity" for hyperbolic sine thus implies that $\sin\beta_i \to 1/\cosh(d_i^{(0)}/2)$ for $i > 0$, and $\sin\beta_0 \to 1/\cosh(H_0/2)$, as $(d_0, d_1, \dots d_{n-1}) \to (H_0, d_1^{(0)}, \dots, d_{n-1}^{(0)})$.

If $(d_0^{(k)}, \dots, d_{n-1}^{(k)})$ converges to $(d_0^{(0)}, \dots, d_{n-1}^{(0)}) \in \sigma^{-j}(\mathrm{Graph}(h_0)$ as $k \to \infty$, then the $\mathbb{Z}_n$-equivariance property of the β_i and the above give:

$$\lim_{k\to\infty} \beta_i(d_0^{(k)}, \dots, d_{n-1}^{(k)}) = \lim_{k\to\infty} \beta_{i-j}(d_j^{(k)}, \dots, d_{n-1}^{(k)}, d_0^{(k)}, \dots, d_{j-1}^{(k)}) = \sin^{-1}\left(\frac{1}{\cosh(d_i^{(0)}/2)}\right)$$

A similar argument shows that α_i has a well-defined extension to the constant 0-function one all of $\widetilde{\mathcal{HC}}_n$, and in particular satisfies $\mathbb{Z}_n$-equivariance. □

The corollary below follows directly from Lemma 4.8 above and the definition in Lemma 3.8, bearing in mind that by Lemma 4.7, $\widetilde{\mathcal{HC}}_n$ has a neighborhood disjoint from $\widetilde{\mathcal{C}}_n$.

Corollary 4.9. *For $n \geq 3$, the functions ν_i extend to* $\mathrm{Graph}(h_0) = U_n \cap \widetilde{\mathcal{HC}}_n$ *by:*

$$\nu_i(d_0, \dots, d_{n-1}) = \begin{cases} \sin^{-1}\left(\frac{1}{\cosh(d_i/2)}\right) + \sin^{-1}\left(\frac{1}{\cosh(d_{i+1}/2)}\right) & i \notin \{0, n-1\} \\ \sin^{-1}\left(\frac{1}{\cosh(d_1/2)}\right) - \sin^{-1}\left(\frac{1}{\cosh(d_0/2)}\right) & i = 0 \\ \sin^{-1}\left(\frac{1}{\cosh(d_{n-1}/2)}\right) - \sin^{-1}\left(\frac{1}{\cosh(d_0/2)}\right) & i = n-1 \end{cases}$$

They extend continuously to all of $\widetilde{\mathcal{HC}}_n$ by the $\mathbb{Z}_n$-equivariance property $\nu_i = \nu_{i-j} \circ \sigma^j$.

The final result of this section has a different flavor; it might best be interpreted as asserting that the subspace topology that $\widetilde{\mathcal{AC}}_n$ inherits from $\mathbb{R}^n$ is geometrically natural.

Proposition 4.10. *Fix $n \geq 3$, $v \in \mathbb{H}^2$, and a geodesic ray δ from v. Each $(d_0, \dots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n$ represents a unique cyclic n-gon (in the sense of Definition 3.1) with center v and $x_0 = \gamma_0 \cap \gamma_1 \in \delta$. A continuous map $\widetilde{\mathcal{AC}}_n \to (\mathbb{H}^2)^n$ is given by $(d_0, \dots, d_{n-1}) \mapsto (x_0, \dots, x_{n-1})$, where $x_i = \gamma_i \cap \gamma_{i+1}$ for each i.*

Proof. For P represented by $(d_0, \dots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n$, since $\mathrm{Isom}^+(\mathbb{H}^2)$ acts transitively there is an isometry that moves P to have its center at v. Then by Lemma 3.6 all vertices of P lie on the circle C of radius $J \doteq J(d_0, \dots, d_{n-1})$ centered at v. The stabilizer of v in $\mathrm{Isom}^+(\mathbb{H}^2)$ acts simply transitively on C, so a unique element of this group takes x_0 to the unique point of $C \cap \delta$. Proposition 2.7 now implies that P is unique with center v and $x_0 \in \delta$ among cyclic n-gons represented by $(d_0, \dots, d_{n-1})$.

Now suppose $(d_0^{(k)},\dots,d_{n-1}^{(k)}) \to (d_0,\dots,d_{n-1})$ as $k\to\infty$, and let cyclic n-gons and $P^{(k)}$ satisfy the criteria of the lemma. For each k let $J^{(k)} = J(d_0^{(k)},\dots,d_{n-1}^{(k)})$, and for $0\le i<n$ let $\alpha_i^{(k)} = \alpha_i(d_0^{(k)},\dots,d_{n-1}^{(k)})$. By Proposition 4.1, J and the α_i are continuous, so $J^{(k)}\to J$ and $\alpha_i^{(k)} \to \alpha_i \doteq \alpha_i(d_0,\dots,d_{n-1})$ for each i, as $k\to\infty$.

For each i let δ_i be the ray from v in the direction of x_i, and let $\delta_i^{(k)}$ be in the direction of $x_i^{(k)}$. Since $\delta_0 = \delta_0^{(k)} = \delta$ by hypothesis, and x_0 and $x_0^{(k)}$ lie at respective distances J and $J^{(k)}$ from v along δ (cf. Lemma 3.6), it follows immediately that $x_0^{(k)}\to x_0$ as $k\to\infty$. We show below that $x_i^{(k)}\to x_i$ as $k\to\infty$ for each i, thus establishing the lemma.

Assume first that $(d_0,\dots,d_{n-1})\in\widetilde{\mathcal{C}}_n$, so $\sum_{i=0}^{n-1}\alpha_i = 2\pi$ by Lemma 3.7. Then $(d_0^{(k)},\dots,d_{n-1}^{(k)})\in\widetilde{\mathcal{C}}_n$ for all but finitely many k, so $\sum_{i=0}^{n-1}\alpha_i^{(k)} = 2\pi$. The angle from δ_0 to δ_i, measured in the direction of δ_1, is $\theta_i = \sum_{j=1}^{i}\alpha_j$ for $i>0$; and the angle from δ_0 to $\delta_i^{(k)}$ in the same direction is $\theta_i^{(k)} = \sum_{j=1}^{i}\alpha_j^{(k)}$. Thus the hyperbolic law of cosines gives:

$$\cosh d(x_i,x_i^{(k)}) = \cosh J^{(k)}\cosh J - \sinh J^{(k)}\sinh J\cos\left|\theta_i-\theta_i^{(k)}\right|$$

Since $J^{(k)}\to J$ and $\left|\theta_i-\theta_i^{(k)}\right|\to 0$ as $k\to\infty$, the above implies that $d(x_i^{(k)},x_i)\to 0$.

For $(d_0,\dots,d_{n-1})\in\widetilde{\mathcal{AC}}_n-\widetilde{\mathcal{C}}_n$ the argument is similar but requires two cases: for U_n as in Lemma 3.2, if $(d_0,\dots,d_{n-1})\in\sigma^{-m}(U_n)$ then one takes $\theta_i = \sum_{j=1}^{i}\alpha_j$ for $0<i\le m$ and $\theta_i = \sum_{j=i}^{n-1}\alpha_j$ for $m<i\le n-1$. For all but finitely many k, $(d_0^{(k)},\dots,d_{n-1}^{(k)})$ is also in $\sigma^{-m}(U_n)$, and defining $\theta_i^{(k)}$ analogously, the argument above goes through.

One additional wrinkle is that for $(d_0,\dots,d_{n-1})\in\widetilde{\mathcal{BC}}_n$ one should pass to two subsequences, one consisting of those $(d_0^{(k)},\dots,d_{n-1}^{(k)})\in\widetilde{\mathcal{C}}_n$ and the other of those in $\widetilde{\mathcal{AC}}_n-\widetilde{\mathcal{C}}_n$, and apply the respective arguments above separately to the relevant subsequences. (Recall from Lemma 3.7 that in this case $\sum_{i=0}^{n-1}\alpha_i = 2\pi$ and $\sum_{i\neq m}\alpha_i = \alpha_m = \pi$.) This finishes the proof. $\square$

5. Defects of packings in cyclic polygons

We now turn our attention to the area of cyclic polygons. The well-known "Gauss-Bonnet formula for hyperbolic polygons" asserts that the area of an n-gon $P\in\mathbb{H}^2$ is equal to its *angle defect*: $(n-2)\pi-\sum_{i=0}^{n-1}\nu_i$, where ν_i is the angle at the ith vertex of P. Here we consider an extended notion of defect, allowing disks of equal radius to be centered at vertices of P.

Definition 5.1. For $n\ge 3$, $(d_0,\dots,d_{n-1})\in\widetilde{\mathcal{AC}}_n$, and $R\ge 0$, let:

$$D_R(d_0,\dots,d_{n-1}) = (n-2)\pi - 2\cosh R\sum_{i=0}^{n-1}\nu_i(d_0,\dots,d_{n-1}),$$

be the *radius-R defect* of $P=(d_0,\dots,d_{n-1})$, where ν_i is as in Lemma 3.8. As with J, we refer by the same name to the function $\mathcal{AC}_n\to\mathbb{R}$ induced by D_R.

That D_R is invariant under the $\mathbb{Z}_n$-action, and hence induces a function on $\mathcal{AC}_n$ follows from the $\mathbb{Z}_n$-equivariance property of the ν_i described in Lemma 3.8. Proposition 4.4 implies that

D_R is C^1 on $\widetilde{\mathcal{AC}}_n$, and our second task in this section is to describe its derivative. First though, we will describe its geometric meaning, for certain R, on certain subsets of $\widetilde{\mathcal{AC}}_n$.

If $U \subset \mathbb{H}^2$ is a disk centered at x, then for $0 \leq \alpha \leq \pi$, a *sector of U with angle measure α* is $U \cap (\mathcal{H} \cap \mathcal{H}')$, where $\mathcal{H}$ and $\mathcal{H}'$ are half-spaces bounded by geodesics γ and γ', respectively, with $x = \gamma \cap \gamma'$ and α the interior angle at x between $\mathcal{H}$ and $\mathcal{H}'$. If U is centered at a vertex x of a polygon P, the *sector of U determined by P* is $U \cap (\mathcal{H} \cap \mathcal{H}')$, where $\mathcal{H}$ and $\mathcal{H}'$ are the half-planes containing P and bounded by geodesics containing the edges of P that intersect at x. It follows that $\angle_v P$ is the angle measure of this sector, which clearly contains $P \cap U$. Containment may be proper in general, but we have:

Lemma 5.2. *Let T be an isosceles triangle with equal sides of length J and a third, γ, with length d. Let $A_d(J)$ and $B_d(J)$ be as in Lemma 1.7, and for $R \geq 0$ define:*

$$D_R(d, J) = \pi - A_d(J) - 2\cosh R \cdot B_d(J),$$

If $R \leq d/2$ and U and U' are disks with radius R centered at the endpoints of γ then $U \cap U' = \emptyset$ and $T \cap U$ (or respectively, U') is the sector of U (or U') determined by T, with angle measure $B_d(J)$. $D_R(d, J)$ is the area of the set of points in T not contained in $U \cup U'$.

Proof. Recall from Lemma 1.7 that $A_d(J)$ is the angle of T at the vertex v opposite γ, and $B_d(J)$ is the angle at each other vertex. Thus by the Gauss–Bonnet formula T has area $\pi - A_d(J) - 2B_d(J)$. By another well known formula, the area of a hyperbolic disk with radius R is $2\pi(\cosh R - 1)$, and a sector with angle measure β has area $\beta(\cosh R - 1)$. We will show that $U \cap P$ and $U' \cap P$ are sectors and hence prove the lemma, since U and U' do not overlap (their centers are at distance $d \geq 2R$), and

$$D_R(d, J) = (\pi - A_d(J) - 2B_d(J)) - 2B_d(J) \cdot (\cosh R - 1).$$

Let δ be the geodesic arc in T from v to its nearest point on γ. Since T is isosceles, δ is an axis of mirror symmetry for T, intersecting γ perpendicularly at its midpoint. Let u be the vertex of γ that is the center of U. Then $d(u, \delta \cap \gamma) = d/2 \geq R$. Each x in $\delta - (\delta \cap \gamma)$ determines a right triangle in T together with u and $\delta \cap \gamma$, and the hyperbolic law of cosines implies

$$\cosh d(u, x) = \cosh d/2 \cosh d_x,$$

where d_x is the distance from x to $\delta \cap \gamma$. In particular, $d(u, x) > d/2 \geq R$ for such x. It follows that the geodesic arc in T from u to x contains the entire intersection between U and the geodesic ray in that direction. Therefore U intersects T in a full sector of angle $B_d(J)$, and since U and U' are interchanged by the mirror symmetry of T, the claim is proved. ☐

A non-centered cyclic polygon P may exhibit the pathology showed in Figure 5.1, in which $P \cap U_1$ is a proper subset of the sector of U_1 determined by P. The lemma below shows that this does not occur in the centered case, and uses the isosceles decomposition to extract a more subtle but still useful conclusion in the non-centered case.



FIGURE 5.1. A pathological packing in the non-centered case.

Lemma 5.3. *For $n \geq 3$ and $(d_0, \ldots, d_{n-1}) \in \overline{U}_n \cap \widetilde{\mathcal{AC}}_n$, where U_n is as in Lemma 3.2, let P be a cyclic n-gon represented by $(d_0, \ldots, d_{n-1})$ (see Definition 3.1), and let $\{T_i\}_{i=0}^{n-1}$ be produced as in Lemma 1.6. If $R \leq \min_i\{d_i/2\}$ and for each i, U_i is an open disk of radius R centered at $u_i = \gamma_i \cap \gamma_{i+1}$ (taking $i+1$ modulo n), then:*

- *If $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$ then $P \cap U_i$ is a sector with angle measure $\nu_i(d_0, \ldots, d_{n-1})$ for each $i \in \{0, \ldots, n-1\}$, and $P \cap U_i \cap U_j = \emptyset$ for each $j \neq i$.*
- *If $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$ then for each $i \in \{1, \ldots, n-2\}$, $(P \cup T_0) \cap U_i$ is a sector with angle measure $\nu_i(d_0, \ldots, d_{n-1})$, $(P \cup T_0) \cap U_0$ is a sector with angle measure $\beta_1(d_0, \ldots, d_{n-1})$, and $(P \cup T_0) \cap U_{n-1}$ is a sector with angle measure $\beta_{n-1}(d_0, \ldots, d_{n-1})$. Furthermore, $(P \cup T_0) \cap U_i \cap U_j = \emptyset$ for $j \neq i$.*

Proof. P as above has radius $J \doteq J(d_0, \ldots, d_{n-1})$ by Lemma 3.6. Let v be the center of P. Since $\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$ is the closure in $\widetilde{\mathcal{AC}}_n$ of $\widetilde{\mathcal{C}}_n$, if it contains $(d_0, \ldots, d_{n-1})$ then Lemma 3.9 implies $v \in P$. Thus by Lemma 1.10, for each i:

$$\angle_{u_i} P = B_{d_i}(J) + B_{d_{i+1}}(J) = \beta_i(d_0, \ldots, d_{n-1}) + \beta_{i+1}(d_0, \ldots, d_{n-1}) = \nu_i(d_0, \ldots, d_{n-1})$$

(Above we have used Lemmas 3.7 and 3.8 to rewrite in terms of the β_i and ν_i.) It follows that $P \cap U_i$ is contained in a sector of U_i with angle measure $\nu_i(d_0, \ldots, d_{n-1})$. Lemma 1.6 implies that T_i and T_{i+1} do not overlap and are contained in P, and Lemma 5.2 implies that they intersect U_i in full sectors with respective angle measures $B_{d_i}(J) = \beta_i(d_0, \ldots, d_{n-1})$ and $B_{d_{i+1}}(J) = \beta_{i+1}(d_0, \ldots, d_{n-1})$. Therefore

$$P \cap U_i = (T_i \cap U_i) \cup (T_{i+1} \cap U_i)$$

is the sector with angle measure $\nu_i(d_0, \ldots, d_{n-1})$.

If $(d_0, \ldots, d_{n-1}) \in \overline{U}_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$, then Lemma 1.10 implies:

$$\angle_{u_0} P = B_{d_1}(J) - B_{d_0}(J) = \beta_1(d_0, \ldots, d_{n-1}) - \beta_0(d_0, \ldots, d_{n-1})$$

Lemma 1.6 implies that $T_1 \cap U_0 \subset (P \cup T_0) \cap U_0$, so since T_0 has angle $\beta_0(d_0, \ldots, d_{n-1})$ at u_0 and Lemma 5.2 implies that $T_1 \cap U_0$ is a full sector, the above implies that $(P \cup T_0) \cap U_0 = T_1 \cap U_0$, a sector with angle measure $\beta_1(d_0, \ldots, d_{n-1})$.

An analogous argument gives the conclusion for $(P \cup T_0) \cap U_{n-1}$ in this case. Again taking $(d_0, \ldots, d_{n-1}) \in \overline{U}_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$, now suppose $1 < i \leq n-1$. Lemma 1.6 implies that T_i and T_{i+1} do not overlap and are contained in $P \cup T_0$, and Lemma 5.2 again implies that they intersect u_i in full sectors with angle measure $B_{d_i}(J) = \beta_i(d_0, \ldots, d_{n-1})$ and $B_{d_{i+1}}(J) = \beta_{i+1}(d_0, \ldots, d_{n-1})$, respectively. Since Lemma 1.10 implies

$$\angle_{u_i} P = \beta_i(d_0, \ldots, d_{n-1}) + \beta_{i+1}(d_0, \ldots, d_{n-1}) = \nu_i(d_0, \ldots, d_{n-1}),$$

it follows that $(P \cup T_0) \cap U_i$ is a full sector with angle measure $\nu_i(d_0, \ldots, d_{n-1})$. □

Lemma 5.4. *For $n \geq 3$ and $(d_0, \ldots, d_{n-1}) \in \overline{U}_n \cap \widetilde{\mathcal{AC}}_n$, let $J = J(d_0, \ldots, d_{n-1})$ from Lemma 3.6. For $R \geq 0$ and $D_R(d, J)$ as in Lemma 5.2:*

$$D_R(d_0, \ldots, d_{n-1}) = \begin{cases} \sum_{i=0}^{n-1} D_R(d_i, J) & (d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n \\ \left(\sum_{i=1}^{n-1} D_R(d_i, J)\right) - D_R(d_0, J) & (d_0 \ldots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n \end{cases}$$

Fix a cyclic n-gon P represented by $(d_0, \ldots, d_{n-1})$, take $R \leq \min_i\{d_i/2\}$, and for each i let U_i be a disk of radius R centered at $u_i = \gamma_i \cap \gamma_{i+1}$. Then:

- *If $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$, then $D_R(d_0, \ldots, d_{n-1})$ is the area of $P - \bigcup_{i=0}^{n-1} U_i$.*
- *If $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$, then $D_R(d_0, \ldots, d_{n-1}) + D_R(d_0, J)$ is the area of $(P \cup T_0) - \bigcup_{i=0}^{n-1} U_i$, where T_0 is as in Lemma 1.6.*

Proof. Since $J = J(d_0, \ldots, d_{n-1})$, $A_{d_i}(J) = \alpha_i(d_0, \ldots, d_{n-1})$ and $B_{d_i}(J) = \beta_i(d_0, \ldots, d_{n-1})$ by Lemma 3.7. Lemma 5.2 therefore gives:

$$D_R(d_i, J) = \pi - \alpha_i(d_0, \ldots, d_{n-1}) - 2\cosh R \cdot \beta_i(d_0, \ldots, d_{n-1})$$

If $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$ then using the identity $2\pi = \sum \alpha_i$ from Lemma 3.7 yields:

$$\begin{aligned} \sum_{i=0}^{n-1} D_R(d_i, J) &= n\pi - \sum_{i=0}^{n-1} \alpha_i(d_0, \ldots, d_{n-1}) - 2\cosh R \sum_{i=0}^{n-1} \beta_i(d_0, \ldots, d_{n-1}) \\ &= (n-2)\pi - \cosh R \sum_{i=0}^{n-1} [\beta_i(d_0, \ldots, d_{n-1}) + \beta_{i+1}(d_0, \ldots, d_{n-1})] \\ &= (n-2)\pi - \cosh R \sum_{i=0}^{n-1} \nu_i(d_0, \ldots, d_{n-1}) = D_R(d_0, \ldots, d_{n-1}) \end{aligned}$$

The final line above uses Lemma 3.8 and Definition 5.1. For $(d_0,\dots,d_{n-1}) \in \overline{U}_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$ the identity $\alpha_0 = \sum \alpha_i$ from Lemma 3.7 gives:

$$\begin{aligned}\left(\sum_{i=1}^{n-1} D_R(d_i,J)\right) - D_R(d_0,J) &= (n-2)\pi + \alpha_0(d_0,\dots,d_{n-1}) - \sum_{i=1}^{n-1} \alpha_i(d_0,\dots,d_{n-1}) \\ &\qquad + 2\cosh R\left[\beta_0(d_0,\dots,d_{n-1}) - \sum_{i=1}^{n-1}\beta_i(d_0,\dots,d_{n-1})\right] \\ &= (n-2)\pi - \cosh R\left[\sum_{i=1}^{n-2}\beta_i(d_0,\dots,d_{n-1}) + \beta_{i+1}(d_0,\dots,d_{n-1})\right. \\ &\qquad \left. - (\beta_1-\beta_0)(d_0,\dots,d_{n-1}) + (\beta_{n-1}-\beta_0)(d_0,\dots,d_{n-1})\right]\end{aligned}$$

This is again equal to $D_R(d_0,\dots,d_{n-1})$ by definition of ν_i.

Recall that if U is a an open hyperbolic disk with radius R, a sector of U with angle measure ν has area $\nu(\cosh R - 1)$. Lemma 5.3 thus directly implies that for $(d_0,\dots,d_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$,

$$D_R(d_0,\dots,d_{n-1}) = \left[(n-2)\pi - \sum_{i=0}^{n-1}\nu_i(d_0,\dots,d_{n-1})\right] - (\cosh R - 1)\sum_{i=0}^{n-1}\nu_i(d_0,\dots,d_{n-1})$$

is the area in P complementary to $P \cap \left(\bigcup_{i=0}^{n-1} U_i\right)$.

For $(d_0,\dots,d_{n-1}) \in \overline{U}_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$ let us recall from Lemma 1.5 that T_0 does not overlap P and $P \cup T_0$ is a convex polygon, and from Lemma 1.7 that T_0 has angles $A_{d_0}(J) = \alpha_0(d_0,\dots,d_{n-1})$ at v and $B_{d_0}(J) = \beta_0(d_0,\dots,d_{n-1})$ at u_0 and u_{n-1}. The Gauss-Bonnet formula and Lemma 5.2 thus imply that the area in $P \cup T_0$ complementary to $\bigcup_{i=0}^{n-1} U_i$ is:

$$\begin{aligned}&\left[(n-2)\pi - \sum_{i=0}^{n-1}\nu_i(d_0,\dots,d_{n-1})\right] + \left[\pi - \alpha_0(d_0,\dots,d_{n-1}) - 2\beta_0(d_0,\dots,d_{n-1})\right] \\ &\qquad - (\cosh R - 1)\left[\sum_{i=1}^{n-2}\nu_i(d_0,\dots,d_{n-1}) + \beta_1(d_0,\dots,d_{n-1}) + \beta_{n-1}(d_0,\dots,d_{n-1})\right] \\ &= D_R(d_0,\dots,d_{n-1}) + \left[\pi - \alpha_0(d_0,\dots,d_{n-1}) - 2\beta_0(d_0,\dots,d_{n-1})\right] \\ &\qquad - 2(\cosh R - 1)\beta_0(d_0,\dots,d_{n-1}) = D_R(d_0,\dots,d_{n-1}) + D_R(d_0,J)\end{aligned}$$

(Recall above that $\nu_0 = \beta_1 - \beta_0$ and $\nu_{n-1} = \beta_{n-1} - \beta_0$ on $\overline{U}_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$.) □

Proposition 5.5. *For* $n \geq 3$ *and* $R \geq 0$, $D_R\colon \widetilde{\mathcal{AC}}_n \to \mathbb{R}^+$ *is* C^1 *and symmetric. At* $(d_0,\dots,d_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$*, for each* $i \in \{0,\dots,n-1\}$ *we have:*

$$\frac{\partial}{\partial d_i} D_R(d_0,\dots,d_{n-1}) = \cosh R\sqrt{\frac{1}{\cosh^2(d_i/2)} - \frac{1}{\cosh^2 J(d_0,\dots,d_{n-1})}}$$

This holds for $i > 0$ *on all of* U_n *(see Lemma 3.2). For* $(d_0,\dots,d_{n-1}) \in U_n \cap (\widetilde{\mathcal{AC}}_n \cup \widetilde{\mathcal{C}}_n)$*:*

$$\frac{\partial}{\partial d_0} D_R(d_0,\dots,d_{n-1}) = -\cosh R\sqrt{\frac{1}{\cosh^2(d_0/2)} - \frac{1}{\cosh^2 J(d_0,\dots,d_{n-1})}}$$

At $(d_0,\dots,d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$, $\frac{\partial}{\partial d_i} D_R(d_0,\dots,d_{n-1}) = 0$.

Proof. We pointed out below its definition that D_R is C^1 (because the ν_i are, see Lemma 4.4) and $\mathbb{Z}_n$-invariant (because the the ν_i are $\mathbb{Z}_n$-equivariant, see Lemma 3.8). Given $(d_0,\dots,d_{n-1}) \in \widetilde{\mathcal{AC}}_n$ and a permutation π, let us assume that $(d_0,\dots,d_{n-1}) \in \overline{U}_n$ and $\pi(0) = 0$. Then taking $J = J(d_{\pi(0)},\dots,d_{\pi(n-1)})$, Lemma 5.4 implies:

$$D_R(d_{\pi(0)},\dots,d_{\pi(n-1)}) = \begin{cases} \sum_{i=0}^{n-1} D_R(d_{\pi(i)},J) & (d_{\pi(0)},\dots,d_{\pi(n-1)}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n \\ \sum_{i=1}^{n-1} D_R(d_{\pi(i)},J) - D_R(d_0,J) & (d_{\pi(0)},\dots,d_{\pi(n-1)}) \notin \widetilde{\mathcal{C}}_n \end{cases}$$

Lemma 3.6 implies that $J = J(d_0,\dots,d_{n-1})$, so $D_R(d_{\pi(0)},\dots,d_{\pi(n-1)}) = D_R(d_0,\dots,d_{n-1})$ since $\widetilde{\mathcal{C}}_n$ and $\widetilde{\mathcal{BC}}_n$ are π-invariant. It thus follows from $\mathbb{Z}_n$-invariance that D_R is symmetric.

Using Lemma 3.8, for $(d_0,\dots,d_{n-1}) \in \widetilde{\mathcal{C}}_n$ and $i \in \{0,\dots,n-1\}$ we obtain:

$$\begin{aligned} \frac{\partial}{\partial d_i} D_R(d_0,\dots,d_{n-1}) &= -\cosh R \sum_{j=0}^{n-1} \left(\frac{\partial \beta_j}{\partial d_i}(d_0,\dots,d_{n-1}) + \frac{\partial \beta_{j+1}}{\partial d_i}(d_0,\dots,d_{n-1}) \right) \\ &= \frac{-\cosh R}{\cosh J} \left[\sum_{j=0}^{n-1} \frac{\partial \alpha_j}{\partial d_i}(d_0,\dots,d_{n-1}) \right] + \cosh R \sqrt{\frac{1}{\cosh^2(d_i/2)} - \frac{1}{\cosh^2 J}} \end{aligned}$$

Above we have used the partial derivative formulas of Lemma 4.3. Applying the relevant case of (4.1.1), the sum of partial derivatives of the α_i vanishes.

At $(d_0,\dots,d_{n-1}) \in U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$, the definition of the ν_i gives:

$$\frac{\partial}{\partial d_i} D_R(d_0,\dots,d_{n-1}) = 2\cosh R \left[\frac{\partial \beta_0}{\partial d_i}(d_0,\dots,d_{n-1}) - \sum_{j=1}^{n-1} \frac{\partial \beta_j}{\partial d_i}(d_0,\dots,d_{n-1}) \right]$$

In this case, Lemma 4.3 yields different results for different i. For $i > 0$ we have:

$$\begin{aligned} \frac{\partial}{\partial d_i} D_R(d_0,\dots,d_{n-1}) = \frac{-\cosh R}{\cosh J} &\left[\frac{\partial \alpha_0}{\partial d_i}(d_0,\dots,d_{n-1}) - \sum_{j=1}^{n-1} \frac{\partial \alpha_j}{\partial d_i}(d_0,\dots,d_{n-1}) \right] \\ &+ \cosh R \sqrt{\frac{1}{\cosh^2(d_i/2)} - \frac{1}{\cosh^2 J}} \end{aligned}$$

Applying the relevant case of (4.1.1), the sum of partials of the α_i again vanishes, and we are left with the identical result to the case above. On the other hand, if $i = 0$ then since the extra term is contributed by $\partial\beta_0/\partial d_0$ its sign is negative (by Lemma 4.3) and after applying (4.1.1) we are left with the opposite value from above.

For $(d_0,\dots,d_{n-1} \in \mathrm{Graph}(b_0)$, $\cosh J(d_0,\dots,d_{n-1}) = d_0/2$ by Lemma 3.9. Using the formulas above this implies that $\partial D_R/\partial d_0$ vanishes here. For $(d_0,\dots,d_{n-1}) \in \sigma^{-i}(\mathrm{Graph}(b_0))$, we have:

$$\frac{\partial}{\partial d_i} D_R(d_0,\dots,d_{n-1}) = \frac{\partial}{\partial d_0} D_R(\sigma^i(d_0,\dots,d_{n-1})) = 0$$

Here we have used the fact that $\sigma(d_0,\dots,d_{n-1}) = (d_i,\dots,d_{n-1},d_0,\dots,d_{i-1})$, and the same holds for its derivative, and the fact that D_R is σ-invariant. □

In this section we wish to ultimately produce a means for bounding defect below using side length bounds. The following lemma describes a "monotonic-convex" property of the closure of centered space that will allow us to produce good bounds for such polygons.

Lemma 5.6. *For $n \geq 3$, suppose $(d_0, \ldots, d_{n-1}) \in \overline{U}_n \cap (\widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n)$ and $(d'_0, \ldots, d'_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$ satisfy $d'_i \geq d_i$ for each i. Let $d = \min_i\{d_i\}$ and $D = \max_i\{d'_i\}$, and define:*

$$d_i(t) = \begin{cases} d_i & d + t \leq d_i \\ d + t & d_i \leq d + t \leq d'_i \\ d'_i & \textit{otherwise} \end{cases} \qquad \textit{for } 0 \leq t \leq D - d$$

This defines a path in $\widetilde{\mathcal{C}}_n$ from $(d_0, \ldots, d_{n-1})$ to $(d'_0, \ldots, d'_{n-1})$ such that $(d_0(t), \ldots, d_{n-1}(t)) \in \mathcal{C}_n$ for each $t \in (0, D - d)$.

Proof. For $0 < t \leq d_0 - d$, $\max_i\{d_i(t)\} = d_0(t) = d_0$ by definition, since $(d_0, \ldots, d_{n-1}) \in \overline{U}_n$ implies that d_0 is maximal among the d_i. Recall from Lemma 1.8 that if $d' > d$ then $A_{d'}(J) > A_d(J)$ for each $J \geq d'/2$. Thus:

$$\sum_{i=1}^{n-1} A_{d_i(t)}(d_0(t)/2) = \sum_{i=1}^{n-1} A_{d_i(t)}(d_0/2) > \sum_{i=1}^{n-1} A_{d_i}(d_0/2) \geq \pi$$

The second inequality above follows from Lemma 3.3 and the hypothesis that $(d_0, \ldots, d_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$. Therefore for $t \leq d_0 - d$, $(d_0(t), \ldots, d_{n-1}(t)) \in \widetilde{\mathcal{C}}_n$ by Definition 3.1.

If $d_0 = D$ then the above case completes the proof, so assume $D > d_0$. Let $i_0 \in \{0, \ldots, n-1\}$ be such that $D = d'_{i_0}$, and let d'_{i_1} be maximal among the d'_i with $i \neq i_0$. Then either $d_0 < d'_{i_1}$ or $d'_{i_1} < d'_{i_0}$.

First suppose that $d_0 < d'_{i_1}$. Then for $d_0 - d \leq t \leq d'_{i_1} - d$, at least $d_{i_0}(t)$ and $d_{i_1}(t)$ take the maximum value $d + t$ among the entries $d_i(t)$. Therefore:

$$\sum_{i \neq i_0} A_{d_i(t)}(d_{i_0}(t)) = \sum_{i \neq i_0} A_{d_i(t)}((d+t)/2) = \pi + \sum_{i \neq i_0, i_1} A_{d_i(t)}((d+t)/2) > \pi$$

Thus $(d_0(t), \ldots, d_{n-1}(t)) \in \mathcal{C}_n$ for $d_0 - d \leq t \leq d'_{i_1} - d$. If $d'_{i_1} = d'_{i_0} = D$ this completes the proof.

If $d_0 \geq d'_{i_1}$, then we skip the case above. Assume $d'_{i_1} < D$; then for $d'_{i_1} - d \leq t < D - d$, $d_{i_0}(t) = d + t$ is the unique maximal element of $\mathcal{D}(t)$, and for $i \neq i_0$, $d_i(t) = d'_i$. Therefore:

$$\sum_{i \neq i_0} A_{d_i(t)}(d_{i_0}(t)/2) = \sum_{i \neq i_0} A_{d'_i}((d+t)/2) > \sum_{i \neq i_0} A_{d'_i}(D/2) \geq \pi$$

The first inequality above follows from the fact that $A_d(t)$ decreases in t for any d, and the second from Lemma 3.3, since $(d'_0, \ldots, d'_{n-1}) \in \widetilde{\mathcal{C}}_n \cup \widetilde{\mathcal{BC}}_n$. Thus $(d_0(t), \ldots, d_{n-1}(t)) \in \widetilde{\mathcal{C}}_n$ for $d'_{i_1} - d \leq t < D - d$. □

Definition 5.7. For $n \geq 3$ and $[d_0, \ldots, d_{n-1}]$ and $[d'_0, \ldots, d'_{n-1}]$ in $\overline{\mathcal{C}}_n$, we say $[d_0, \ldots, d_{n-1}] < [d'_0, \ldots, d'_{n-1}]$ if there exists j such that $d_{\sigma^j(i)} \leq d'_i$ for all i and for some i_0, $d_{\sigma^j(i_0)} < d'_{i_0}$.

Corollary 5.8. *For $n \geq 3$, if $[d_0, \ldots, d_{n-1}] < [d'_0, \ldots, d'_{n-1}] \in \mathcal{C}_n$, then $D_R[d_0, \ldots, d_{n-1}] < D_R[d'_0, \ldots, d'_{n-1}]$.*

Proof. Given elements $[d_0, \ldots, d_{n-1}] \leq [d'_0, \ldots, d'_{n-1}]$ of $\overline{\mathcal{C}}_n$, by Definition 5.7 we may choose pre-images $(d_0, \ldots, d_{n-1})$ and $(d'_0, \ldots, d'_{n-1})$ in $\widetilde{\mathcal{C}}_n$ with $d_i \leq d'_i$ for each i; and furthermore, after applying the same power of σ to each, we may take $(d_0, \ldots, d_{n-1}) \in \overline{U}_n$. Lemma 5.6 thus produces a path $(d_0(t), \ldots, d_{n-1}(t))$ from $(d_0, \ldots, d_{n-1})$ to $(d'_0, \ldots, d'_{n-1})$ in $\widetilde{\mathcal{C}}_n$.

Observe that $t \mapsto (d_0(t), \ldots, d_{n-1}(t))$ is piecewise-smooth. For $i \in \{0, \ldots, n-1\}$, computing a derivative from the right yields:

$$\frac{d}{dt} d_i(t) = \begin{cases} 1 & d_i - d \leq t < d'_i - d \\ 0 & \text{otherwise} \end{cases}$$

With d and D as in Lemma 5.6, for $0 \leq t < D - d$ the chain rule yields:

$$\frac{d}{dt} D_R(d_0(t), \ldots, d_{n-1}(t)) = \sum_{i=0}^{n-1} \frac{d}{dt} d_i(t) \cdot \frac{\partial}{\partial d_i} D_R(d_0(t), \ldots, d_{n-1}(t))$$

Here again we compute the derivative from the right. By Proposition 5.5 and Lemma 5.6, the sum above is non-negative for all $t \in [0, D-d)$. It is positive for $t \in (d_{i_0} - d, d'_{i_0} - d)$, where i_0 is such that $d_{i_0} < d'_{i_0}$. □

The lemma below extends the defect function continuously to $\widetilde{\mathcal{AC}}_n \cup \widetilde{\mathcal{HC}}_n$. It is a direct consequence of Definition 5.1 and Corollary 4.9, which extends the vertex angle functions.

Lemma 5.9. *For $n \geq 3$, $R \geq 0$, and $i \in \{0, \ldots, n-1\}$, D_R extends continuously to $\sigma^{-i}(\mathrm{Graph}(h_0))$ by:*

$$D_R(d_0, \ldots, d_{n-1}) = (n-2)\pi + 2\cosh R \left[\sin^{-1}\left(\frac{1}{\cosh(d_i/2)}\right) - \sum_{j \neq i} \sin^{-1}\left(\frac{1}{\cosh(d_j/2)}\right) \right]$$

This extension is $\mathbb{Z}_n$-invariant on $\widetilde{\mathcal{HC}}_n$.

Lemma 5.10. *For $n \geq 3$, suppose that distinct $(d_1, \ldots, d_{n-1})$ and $(d'_1, \ldots, d'_{n-1})$ in $(R^+)^{n-1}$ satisfy $d'_i \geq d_i$ for each i. For $h_0 \colon (\mathbb{R}^+)^{n-1} \to \mathbb{R}^+$ as in Lemma 3.10 and $R \geq 0$,*

$$D_R(h_0(d_1, \ldots, d_{n-1}), d_1, \ldots, d_{n-1}) < D_R(h_0(d'_1, \ldots, d'_{n-1}), d'_1, \ldots, d'_{n-1})$$

Proof. The formula from Lemma 5.9 determines a smooth function $\overline{D}_R$ on a sufficiently small neighborhood V of $\widetilde{\mathcal{HC}}_n$ in $\mathbb{R}^n$. Lemma 3.10 thus implies that f below is smooth on $(\mathbb{R}^+)^{n-1}$:

$$f(d_1, \ldots, d_{n-1}) = D_R(h_0(d_1, \ldots, d_{n-1}), d_1, \ldots, d_{n-1})$$

For fixed $(d_1, \ldots, d_{n-1}) \in (\mathbb{R}^+)^{n-1}$, if $H_0 = h_0(d_1, \ldots, d_{n-1})$ the chain rule gives:

$$\begin{aligned} \frac{\partial f}{\partial d_i}(H_0, d_1, \ldots, d_{n-1}) &= \frac{\partial \overline{D}_R}{\partial d_0}(H_0, d_1, \ldots, d_{n-1}) \frac{\partial h_0}{\partial d_i}(d_1, \ldots, d_{n-1}) + \frac{\partial \overline{D}_R}{\partial d_i}(H_0, d_1, \ldots, d_{n-1}) \\ &= \cosh R \left[\frac{-1}{\cosh(H_0/2)} \cdot \frac{\cosh(d_i/2)}{\cosh(H_0/2)} + \frac{1}{\cosh(d_i/2)} \right] \end{aligned}$$

The second line follows from a computation. Since $H_0 > d_i$ for each $i > 0$ (see the definition in Lemma 3.10), the quantity above is positive. The straight-line path in $(\mathbb{R}^+)^{n-1}$ from $(d_0, \ldots, d_{n-1})$ to $(d'_0, \ldots, d'_{n-1})$, with $d'_i \geq d_i$ for each i, thus determines a path in $\mathrm{Graph}(h_0)$ from $(h_0(d_1, \ldots, d_{n-1}), d_1, \ldots, d_{n-1})$ to $(h_0(d'_1, \ldots, d'_{n-1}), d'_1, \ldots, d'_{n-1})$ along which defect increases. □

Corollary 5.11. *Suppose* $(d'_0, d'_1, \ldots, d'_{n-1}) \in U_n \cap (\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n)$*, and that* $(d_1, \ldots, d_{n-1}) \in (\mathbb{R}^+)^{n-1}$ *satisfies* $d_i \leq d'_i$ *for each* i*. Then with* $H_0 = h_0(d_1, \ldots, d_{n-1})$ *as in Lemma 3.10,* $D_R(d'_0, \ldots .d'_{n-1}) > D_R(H_0, d_1, \ldots, d_{n-1})$ *for* $R \geq 0$*.*

Proof. Let $H'_0 = h_0(d'_1, \ldots, d'_{n-1})$. The path $t \mapsto (t, d'_1, \ldots, d'_{n-1})$, for $t \in [d'_0, H'_0]$, takes $(d'_0, \ldots, d'_{n-1})$ to $(H'_0, d'_1, \ldots, d'_{n-1})$ through $\widetilde{\mathcal{AC}}_n - \widetilde{\mathcal{C}}_n$ (this is easily checked with Definition 3.1). Proposition 5.5 implies that defect strictly decreases along this path for $t < H'_0$, so $D_R(d_0,', \ldots, d'_{n-1}) > D_R(H'_0, d'_1, \ldots, d'_{n-1})$. Lemma 5.10 implies that $D_R(H'_0, d'_1, \ldots, d'_{n-1}) \geq D_R(H_0, d_1, \ldots, d_{n-1})$, and the conclusion follows. □

6. Some explicit formulas

In this section we give explicit formulas, in some important special cases, for several functions that until now have only been defined implicitly; in particular, for the radius function J from Lemma 3.6 on $\widetilde{\mathcal{AC}}_3$ and $\widetilde{\mathcal{AC}}_4$, and for the function b_0 from Lemma 3.4 on $(\mathbb{R}^+)^2$ and $(\mathbb{R}^+)^3$. We also exploit invariance and equivariance properties to describe values at "highly symmetric" points of $\widetilde{\mathcal{AC}}_n$ for arbitrary n. The final result of the section has a different flavor: it minimizes defect over all cyclic triangles that satisfy certain constraints.

Lemma 6.1. *For* $(d_0, d_1, d_2) \subset \widetilde{\mathcal{AC}}_3$ *and* $i \in \{0, 1, 2\}$*, taking* $i \pm 1$ *modulo* 3*:*

$$\cos \nu_i(d_0, d_1, d_2) = \frac{\cosh d_i \cosh d_{i+1} - \cosh d_{i-1}}{\sinh d_i \sinh d_{i+1}}$$

Taking $s_i = \sinh(d_i/2)$ *for each* $i \in \{0, 1, 2\}$*:*

$$\sinh J(d_0, d_1, d_2) = \frac{2 s_0 s_1 s_2}{\sqrt{(s_0 + s_1 + s_2) \prod_{i=0}^{2} (-s_i + \sum_{j \neq i} s_j)}}$$

Proof. Recall from Lemma 3.8 that for $i \in \{0, 1, 2\}$, $\nu_i(d_0, d_1, d_2)$ is the angle of a cyclic triangle P with cyclically ordered side length collection (d_0, d_1, d_2), at the intersection of the sides with lengths d_i and d_{i+1}. The hyperbolic law of cosines thus gives the above formula for $\nu_i(d_0, d_1, d_2)$.

Since J is $\mathbb{Z}_3$-invariant, we may assume (d_0, d_1, d_2) is in the fundamental domain $\overline{U}_3$ for the action (see Lemma 3.2). Then either $\alpha_0(d_0, d_1, d_2)$ or $2\pi - \alpha_0(d_0, d_1, d_2)$ is equal to $\alpha_1(d_0, d_1, d_2) + \alpha_2(d_0, d_1, d_2)$ by Lemma 3.7. Since $\cos(2\pi - \alpha_0) = \cos \alpha_0$, the angle addition formula for cosine yields the following formula relating the α_i in both the centered and non-centered cases:

$$\cos \alpha_0 = \cos \alpha_1 \cos \alpha_2 - \sin \alpha_1 \sin \alpha_2$$

The formula for $J = J(d_0, d_1, d_2)$ is obtained by substituting $A_{d_i}(J)$ for α_i above, using the description from Lemma 1.7, and solving for $\sinh^2 J$. $\square$

Lemma 6.2. *For $(d_1, d_2) \in (\mathbb{R}^+)^2$, $B_0 = b_0(d_1, d_2)$ from Lemma 3.4 satisfies:*

$$\cosh B_0 = \cosh d_1 + \cosh d_2 - 1 \qquad \sinh(B_0/2) = \sqrt{\sinh^2(d_1/2) + \sinh^2(d_2/2)}$$

Proof. Lemma 3.9 implies that $(d_0, d_1, d_2) \in \overline{U}_3 \cap \widetilde{\mathcal{BC}}_3$ if and only if $J(d_0, d_1, d_2) = d_0/2$. Thus by Lemma 3.4, for each $(d_1, d_2) \in (\mathbb{R}^+)^2$, $B_0 = b_0(d_1, d_2)$ uniquely solves $\sinh(B_0/2) = \sinh J(B_0, d_1, d_2)$. Inserting the formula from Lemma 6.1 yields a quadratic equation in $\sinh^2(B_0/2)$, with unique solution $\sinh^2(B_0/2) = \sinh^2(d_1/2) + \sinh^2(d_2/2)$. The left-hand formula above follows using "half-angle" identities. $\square$

We use the explicit formula for J on $\widetilde{\mathcal{AC}}_3$ to give one on $\widetilde{\mathcal{AC}}_4$:

Lemma 6.3. *For $(d_0, d_1, d_2, d_3) \in \widetilde{\mathcal{AC}}_4$, taking $s_i = \sinh(d_i/2)$ for each i:*

$$\sinh J(d_0, d_1, d_2, d_3) = 2\sqrt{\frac{(s_0 s_1 + s_2 s_3)(s_0 s_2 + s_1 s_3)(s_0 s_3 + s_1 s_2)}{\prod_{i=0}^{3}\left(-s_i + \sum_{j\neq i} s_j\right)}}$$

Proof. Let P be a cyclic quadrilateral with cyclically ordered side collection $\gamma_0, \gamma_1, \gamma_2, \gamma_3$ associated to (d_0, d_1, d_2, d_3). A diagonal joining $\gamma_0 \cap \gamma_3$ to $\gamma_1 \cap \gamma_2$ divides P into cyclic triangles T_0 and T_1. If this diagonal has length D then T_0 (say) has cyclically ordered side length collection (d_0, d_1, D) and T_1 has (d_2, d_3, D). We will assume without loss of generality that $\sinh^2(d_0/2) + \sinh^2(d_1/2) \geq \sinh^2(d_2/2) + \sinh^2(d_3/2)$. We have:

$$J(d_0, d_1, D) = J(d_0, d_1, d_2, d_3) = J(d_2, d_3, D)$$

Lemma 6.1 gives formulas for the outermost quantities above, and setting these equal to each other and simplifying yields a quadratic polynomial in $\sinh^2(D/2)$ with coefficients among the $\sinh(d_i/2)$ for $i \in \{0, 1, 2, 3\}$. Using our assumption above, the quadratic formula gives:

$$\sinh^2(D/2) = \frac{s_2 s_3(s_0^2 + s_1^2) + s_0 s_1(s_2^2 + s_3^2)}{s_2 s_3 + s_0 s_1} \quad \text{or} \quad \frac{s_2 s_3(s_0^2 + s_1^2) - s_0 s_1(s_2^2 + s_3^2)}{s_2 s_3 - s_0 s_1} \tag{6.3.1}$$

Our assumption implies that the right-hand quantity is greater than $\sinh^2(d_0/2) + \sinh^2(d_1/2)$ if it is not equal to the left-hand quantity. We claim that $\sinh^2(D/2) \leq \sinh^2(d_0/2) + \sinh^2(d_1/2)$ in all cases, so it is given by the left-hand quantity above.

If P is centered then its center is in one of *int* T_0, *int* T_1, or the diagonal $T_0 \cap T_1$. In the first case, T_0 is centered and T_1 is not, this is reversed in the second case, and in the third case neither is centered but each contains the center. Re-interpreting the criterion of Lemma 6.2 in terms of the hyperbolic sine produces the following inequalities in these cases:

$$s_0^2 + s_1^2 > \sinh^2(D/2) > s_2^2 + s_3^2$$
$$s_0^2 + s_1^2 < \sinh^2(D/2) < s_2^2 + s_3^2$$
$$s_0^2 + s_1^2 = \sinh^2(D/2) = s_2^2 + s_3^2$$

The second case violates our assumption, and in the third the two possibilities from (6.3.1) are equal. We conclude that if P is centered then the claim holds.

If P is non-centered then by Lemma 1.5 a single side separates its center from its interior. If this side is γ_0 or γ_1, say γ_0, then Lemma 6.2 implies that $s_0^2 \geq s_1^2 + \sinh^2(D/2)$, and the claim follows in this case. If the separating side is γ_3 or γ_4, say γ_3, then $s_3^2 \geq s_4^2 + \sinh^2(D/2)$. But since the diagonal of P also separates its center from the interior of $int\, T_0$ we have $\sinh^2(D/2) \geq s_0^2 + s_1^2$ in this case as well, contradicting our assumption that $s_0^2 + s_1^2 \geq s_2^2 + s_3^2$. Thus the claim holds.

Substituting the formula for $\sinh(D/2)$ from (6.3.1) into the formula for $J(d_0, d_1, D)$ gives the result. □

Lemma 6.4. *For* $(d_1, d_2, d_3) \in (\mathbb{R}^+)^3$, $B_0 = b_0(d_1, d_2, d_3)$ *from Lemma 3.4 satisfies:*

$$\sinh(B_0/2) = \left(s_1 s_2 s_3 + \sqrt{(s_1 s_2 s_3)^2 - \left(\frac{s_1^2 + s_2^2 + s_3^2}{3}\right)^3} \right)^{1/3} + \left(s_1 s_2 s_3 - \sqrt{(s_1 s_2 s_3)^2 - \left(\frac{s_1^2 + s_2^2 + s_3^2}{3}\right)^3} \right)^{1/3}$$

Above, $s_i = \sinh(d_i/2)$ *for* $i = 1, 2, 3$, *and the complex cube root is chosen to take the value* 1 *at* 1, *with a branch cut along the negative real axis.*

Proof. As in the proof of Lemma 6.2, it follows from Lemmas 3.9 and 3.4 that $(B_0, d_1, d_2) \in \widetilde{\mathcal{BC}}_n$ if and only if $\sinh(B_0/2) = \sinh J(B_0, d_1, d_2, d_3)$. Substituting the formula from Lemma 6.3 for $J(B_0, d_1, d_2, d_3)$ and moving all terms to one side yields the following equation (after simplification):

$$\left(\sinh^3(B_0/2) - (s_1^2 + s_2^2 + s_3^2)\sinh(B_0/2) - 2s_1 s_2 s_3\right)^2 = 0$$

The left-hand side above is the square of a cubic equation in $\sinh(B_0/2)$. The discriminant D of this cubic is below; the final inequality uses the arithmetic-geometric mean inequality:

$$D = 4(s_1^2 + s_2^2 + s_3^2)^3 - 27(2s_1 s_2 s_3)^2 = 4 \cdot 27 \left[\left(\frac{s_1^2 + s_2^2 + s_3^2}{3}\right)^3 - s_1^2 s_2^2 s_3^2 \right] \geq 0$$

There are thus three real roots, given by Cardano's formula. For s_1, s_2, and s_3 positive, the unique positive root is recorded above. □

Remark 6.5. We will not give an explicit general formula for the radius of a cyclic n-gon for $n > 4$, but the proof of Lemma 6.3 suggests a way of computing numerical values: given a cyclic n-gon, divide it into $n - 2$ cyclic triangles using diagonals from a fixed vertex. Setting the radii of these triangles equal, Lemma 6.1 yields $n - 3$ equations in the diagonal lengths. Solving for the diagonal lengths and inserting into a radius formula gives the result.

Also note that given the radius of a cyclic n-gon, its radius-R defect is easily computed using Lemma 5.4. In highly symmetric cases, the radius and defect can be explicitly described.

Lemma 6.6. *For $d > 0$ and $n \geq 3$, $(d, \dots, d) \in \widetilde{\mathcal{C}}_n$. Furthermore, $J_n(d) \doteq J(d, \dots, d)$ and, for $R \geq 0$, $D_{R,n}(d) \doteq D_R(d, \dots, d)$ satisfy:*

$$\sinh J_n(d) = \sqrt{\frac{2\sinh^2(d/2)}{1-\cos(2\pi/n)}} = \frac{\sinh(d/2)}{\sin(\pi/n)}$$

$$D_{R,n}(d) = (n-2)\pi - 2n\cosh R \sin^{-1}\left(\frac{\cos(\pi/n)}{\cosh(d/2)}\right)$$

For fixed d, $J_n(d)$ increases with n, and the same holds for $D_{R,n}(d)$ when $R \leq d/2$.

Proof. Checking Definition 3.1 gives $(d, \dots, d) \in \widetilde{\mathcal{C}}_n$. This is a fixed point for the $\mathbb{Z}_n$-action on $\mathbb{R}^n$, so it follows from $\mathbb{Z}_n$-equivariance that $\alpha_i(d, \dots, d) = \alpha_j(d, \dots, d)$ for all i and j (see Lemma 3.7). Since $\sum_{i=0}^{n-1} \alpha_i = 2\pi$ we obtain $\alpha_i(d, \dots, d) = 2\pi/n$ for each i. By definition, $\alpha_i(d, \dots, d) = A_d(J_n(d))$, where $A_d(J)$ is as defined in Lemma 1.7, and appealing to the formula there and the fact that $\alpha_i(d, \dots, d) = 2\pi/n$ yields the formula claimed for $J_n(d)$.

Appealing to the definitions in Lemmas 3.7 and 1.7, we find that for any i,

$$\beta_i(d, \dots, d) = \cos^{-1}\left(\frac{\tanh(d/2)}{\tanh J_n(d)}\right) = \cos^{-1}\sqrt{1 - \frac{\cos^2(\pi/n)}{\cosh^2(d/2)}} = \sin^{-1}\left(\frac{\cos(\pi/n)}{\cosh(d/2)}\right)$$

Here we have used trigonometric and hyperbolic-trigonometric identities to simplify formulas, in particular applying the identity $\sin^{-1} x = \cos^{-1}\sqrt{1-x^2}$ for the final equation. The vertex angle function $\nu_i(d, \dots, d) = \beta_i(d, \dots, d) + \beta_{i+1}(d, \dots, d)$ for each i, by Lemma 3.8, so Definition 5.1 yields the formula for $D_{R,n}(d, \dots, d)$ claimed here, using the description of the β_i derived above.

It is easy to see directly that $J_n(d)$ increases with n. For $D_{R,n}(d)$ we consider the function $D_{R,n}(d)/n$. Using the formula for $D_{R,n}(d)$ and taking a derivative with respect to n yields:

$$\frac{\partial}{\partial n}\left(\frac{D_{R,n}(d)}{n}\right) = \frac{2\pi}{n^2}\left(1 - \frac{\cosh R}{\cosh(d/2)} \frac{\sin(\pi/n)}{\sqrt{\cosh^2(d/2) - \cos^2(\pi/n)}}\right)$$

For $R \leq d/2$ this quantity is positive, and the final claim follows. □

Lemma 6.7. *For $R \geq 0$, $d > 0$, and $n \geq 3$, with $b_0 \colon (\mathbb{R}^+)^{n-1} \to \mathbb{R}$ as in Lemma 3.4 and $J_{2(n-1)}(d)$ and $D_{R,2(n-1)}(d)$ as in Lemma 6.6, $B_0 \doteq b_0(d, \dots, d)$ satisfies:*

$$J(B_0, d, \dots, d) = J_{2(n-1)}(d) \qquad D_R(B_0, d, \dots, d) = \frac{1}{2} D_{R,2(n-1)}(d),$$

For fixed d and $R \geq d/2$, these quantities increase with n. Furthermore:

$$\cosh B_0 = \frac{2\cosh d - 2}{1 - \cos(\pi/(n-1))} + 1$$

Proof. Let P be a cyclic n-gon with cyclically ordered side length collection $(b_0, d, \dots, d)$. By Lemma 3.9, the center of P is the midpoint of the side γ_0 with length b_0. Doubling P across γ_0 thus yields a centered $2(n-1)$-gon Q with all sides of length d and the same center and radius as P. Lemma 6.6 thus implies that $J(b_0, d, \dots, d) = J_{2(n-1)}(d)$ (recall from Lemma 3.6 that J assigns to a tuple the radius of the corresponding cyclic polygon).

For each vertex of x of P not contained in γ_0, Q has two vertices with angle measure $\angle_x P$. For each endpoint $x_\pm$ of γ_0, Q has a single vertex with angle measure $2\angle_{x_\pm} P$. That $D_R(b_0, d, \dots, d) = \frac{1}{2} D_{R,2(n-1)}(d)$ thus follows from the formula in Definition 5.1 and the fact that the functions ν_i assign to a tuple the angle of the corresponding cyclic polygon at its ith vertex (see Lemma 3.8).

The value of B_0 recorded above follows by setting $B_0 = 2J_{2(n-1)}(d)$ (see Lemma 3.4). $\square$

Lemma 6.8. *For $d, d' > 0$, (d, d', d, d') and (d, d', d', d) are in $\widetilde{\mathcal{C}}_4$. For $R \geq 0$,*

$$J(d, d', d, d') = J(d, d', d', d) = B_0/2$$
$$D_R(d, d', d, d') = D_R(d, d', d', d) = 2D_R(d, d', B_0),$$

where $B_0 = \cosh^{-1}(\cosh d + \cosh d' - 1) = b_0(d, d')$, for $b_0 \colon (\mathbb{R}^+)^2 \to \mathbb{R}$ as in Lemma 3.4. A centered quadrilateral with cyclically ordered side length collection (d, d', d, d') or (d, d', d', d) has a diagonal that contains its center and divides it into isometric triangles, each with cyclically ordered side length collection S_3-equivalent to $(B_0, d, d') \in \widetilde{\mathcal{BC}}_3$.

Proof. We may assume without loss of generality, by cyclically rearranging if necessary, that $d \geq d'$. For either (d, d', d, d') or (d, d', d', d), $A_d(d/2) = \pi$ occurs twice in the sum from Definition 3.1 that diagnoses centeredness, so each is in $\widetilde{\mathcal{C}}_4$.

Let P and P' be centered quadrilaterals with cyclically ordered side length collections (d, d', d, d') and (d, d', d', d), respectively. Note that (d, d', d, d') is fixed by the order-2 element of $\mathbb{Z}_4$, and (d, d', d', d) is fixed by $\tau \colon (d_0, d_1, d_2, d_3) \mapsto (d_3, d_2, d_1, d_0)$. Therefore by Proposition 2.7, each of P and P' has an order-2 isometry, which preserves orientation in the case of P and reverses it in the case of P'.

The isometry of P exchanges opposite vertices in pairs, so it preserves a geodesic arc γ joining an arbitrarily chosen vertex to its opposite. The isometry of P' fixes two opposite vertices, and exchanges the other two, so in particular it fixes an arc γ' joining the two fixed vertices. Each isometry fixes the center of the corresponding quadrilateral but exchanges complementary components to the arc identified above. It follows that γ contains the center of P, and γ' contains the center of P'.

Let B be the length of γ. P is divided by γ into two cyclic triangles, each with cyclically ordered side length collection (B, d, d') (or (B, d', d) depending on the choice of endpoints of γ). Since each has its center in the edge γ, Lemma 3.9 implies that $(B, d, d') \in \widetilde{\mathcal{BC}}_3$. Hence $B = b_0(d, d')$ by Lemma 3.4, so Lemma 6.2 implies that $B = B_0$ as defined above. It similarly follows that γ' has length B_0: the only difference is that it divides P' into one triangle with cyclically ordered side length collection (B, d, d') and one with (B, d', d), but this does not alter the argument.

P has radius equal to that of its constituent triangles, and so does P', so Lemma 3.9 implies that $J(d, d', d, d') = J(d, d', d', d) = B_0/2$. If a triangle with cyclically ordered side length collection (B_0, d, d') has angles μ_0 and μ_1 at the endpoints of the side γ with length B_0, and angle ν at the vertex opposite γ, then P has two vertices with angle measure ν and two with measure $\mu_0 + \mu_1$. P' has angle ν at two vertices, and one each with angle 2α and 2β. The defect assertion thus follows from Definition 5.1 and Lemma 3.8. □

Lemma 6.9. *Fix $d_0 > 0$, $R \leq d_0/2$, and B_0 such that $\cosh B_0 \geq 2\cosh d_0 - 1$. Then among all $(B_0, d, d') \in \mathcal{BC}_3$ with d and d' at least d_0, $D_R(B_0, d, d') \geq D_R(B_0, d_0, d_0')$, where $d_0' = \cosh^{-1}(\cosh B_0 - \cosh d_0 + 1)$.*

Proof. Let $d_1(t)$ satisfy $\cosh d_1(t) = \cosh d_0 + t$, and let $d_2(t)$ satisfy $\cosh d_2(t) = \cosh B_0 - \cosh d_1(t) + 1$, for $t \in [0, \cosh B_0 - (2\cosh d_0 - 1)]$. Then $d_2(t) \geq d_0$ for each t in this interval, and $(B_0, d_1(t), d_2(t)) \in \mathcal{BC}_3$ by Lemma 6.2. Indeed, Lemmas 3.4 and 6.2 imply that each $(B_0, d, d') \in \mathcal{BC}_3$ with d and d' at least d_0 is of the form $(B_0, d_1(t), d_2(t))$ for some t in $[0, \cosh B_0 - (2\cosh d_0 - 1)]$. Proposition 5.5 gives:

$$\begin{aligned}\frac{d}{dt} D_R(B_0, d_1(t), d_2(t)) &= d_1'(t) \cdot \frac{\partial}{\partial d_1} D_R(B_0, d_1(t), d_2(t)) + d_2'(t) \cdot \frac{\partial}{\partial d_2} D_R(B_0, d_1(t), d_2(t)) \\ &= \frac{1}{\sinh d_1(t)} \sqrt{\frac{1}{\cosh^2(d_1(t)/2)} - \frac{1}{\cosh^2(b/2)}} \\ &\qquad - \frac{1}{\sinh d_2(t)} \sqrt{\frac{1}{\cosh^2(d_2(t)/2)} - \frac{1}{\cosh^2(b/2)}}\end{aligned}$$

Thus $D_R(B_0, d_1(t), d_2(t))$ increases with t for as long as $d_1(t) < d_2(t)$, and decreases for $d_1(t) > d_2(t)$. It therefore finds its global minimum at an endpoint of the interval.

We have $d_1(0) = d_0$ and $\cosh d_2(0) = \cosh B_0 - \cosh d_0 + 1$, and for $T = \cosh B_0 - (2\cosh d_0 - 1)$ we have $\cosh d_1(T) = \cosh B_0 - \cosh d_0 + 1 = \cosh d_2(0)$ and $\cosh d_2(T) = \cosh d_0 = \cosh d_1(0)$. Since the map $\tau\colon (d_0, d_1, d_2) \mapsto (d_2, d_1, d_0)$ takes $(B_0, d_1(0), d_2(0))$ to a cyclic permutation of $(B_0, d_1(T), d_2(T)) = (B_0, d_2(0), d_1(0))$, D_R takes the same value on these. Therefore its minimum occurs at both endpoints of $[0, \cosh B_0 - (2\cosh d_0 - 1)]$. □

REFERENCES

[1] Jean-Marc Schlenker. Small deformations of polygons and polyhedra. *Trans. Amer. Math. Soc.*, 359(5):2155–2189, 2007.

[2] Rolf Walter. Polygons in hyperbolic geometry 1: Rigidity and inversion of the n-inequality. Preprint. arXiv:1008.3404v1, August 2010.

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY

E-mail address: `jdeblois@math.stanford.edu`